

# A note on group representations, determinantal hypersurfaces and their quantizations

Igor Klep and Jurij Volčič

**Abstract.** Recently, there have been exciting developments on the interplay between representation theory of finite groups and determinantal hypersurfaces. For example, a finite Coxeter group is determined by the determinantal hypersurface described by its natural generators under the regular representation. This short note solves three problems about extending this result in the negative. On the affirmative side, it is shown that a quantization of a determinantal hypersurface, the so-called free locus, correlates well with representation theory. If  $A_1, \dots, A_\ell \in \mathrm{GL}_d(\mathbb{C})$  generate a finite group  $G$ , then the family of hypersurfaces  $\{X \in M_n(\mathbb{C})^d: \det(I + A_1 \otimes X_1 + \dots + A_\ell \otimes X_\ell) = 0\}$  for  $n \in \mathbb{N}$  determines  $G$  up to isomorphism.

**Mathematics Subject Classification (2010).** Primary 20C15, 15A22; Secondary 47A56, 14J70.

**Keywords.** Linear pencil, group representation, determinantal hypersurface, free locus.

## 1. Introduction

To  $A_0, \dots, A_\ell \in M_d(\mathbb{C})$  one assigns the **determinantal hypersurface**

$$\{[\xi_0 : \dots : \xi_\ell] \in \mathbb{C}\mathbb{P}^\ell: \det(\xi_0 A_0 + \dots + \xi_\ell A_\ell) = 0\}. \quad (1.1)$$

This is a classical object in algebraic geometry [Dic21, dCEP18, Bea00, Dol12], where a key question asks which hypersurfaces admit determinantal representations. When  $A_j$  are real symmetric matrices, determinantal hypersurfaces pertain to hyperbolic and stable polynomials [HV07, Brä11, KV12, Wag11, Vol19]. The geometry of the hypersurface (1.1) is also explored in multivariate operator theory [Yan09, CY13, CSZ16]. If  $A_j$  are bounded operators on a Hilbert space and the determinant in (1.1) is replaced with the

---

The first author was supported by the Slovenian Research Agency grants J1-8132, N1-0057, P1-0222, and partially supported by the Marsden Fund Council of the Royal Society of New Zealand.

condition that  $\xi_0 A_0 + \dots + \xi_\ell A_\ell$  is not invertible, then (1.1) is known as the **projective joint spectrum** of  $A_0, \dots, A_\ell$  (cf. Taylor spectrum [Tay70] for ensembles of commuting operators).

Through the work of Frobenius [Fro68] and Dedekind [Ded68] on group determinants (see also [Dic02]), determinantal hypersurfaces also pertain to representation theory. Several fascinating developments in this direction [ST15, GY17, ČST18] have been recently made. This note addresses certain limitations for extensions of these results.

Let  $G$  be a finitely generated group. If  $T = (g_1, \dots, g_\ell)$  is a finite sequence of generators for  $G$  and  $\rho : G \rightarrow \mathrm{GL}_d(\mathbb{C})$  is a representation of  $G$ , then denote

$$\mathcal{Z}_1(T, \rho) = \{ \xi \in \mathbb{C}^\ell : \det(I_d + \xi_1 \rho(g_1) + \dots + \xi_\ell \rho(g_\ell)) = 0 \}. \quad (1.2)$$

It is natural to ask what kind of information the affine hypersurface  $\mathcal{Z}_1(T, \rho)$  carries about  $\rho$  and  $G$ . For example, if  $G_1, G_2$  are finite groups with left regular representations  $\lambda_1, \lambda_2$ , then  $\mathcal{Z}_1(G_1 \setminus \{1\}, \lambda_1) = \mathcal{Z}_1(G_2 \setminus \{1\}, \lambda_2)$  implies that  $G_1, G_2$  are isomorphic [FS91]. However, one is typically interested in smaller generating sets or in finitely generated groups which are not necessarily finite. In [GY17], the authors computed the joint spectrum for the infinite dihedral group

$$D_\infty = \langle a, t \mid a^2 = t^2 = 1 \rangle$$

with respect to the generating set  $(1, a, t)$ , and analyzed its properties through the representation theory of  $D_\infty$ . Determinantal hypersurfaces also have a strong connection with representation theory in the case of finite Coxeter groups [ČST18]. A Coxeter group is a finitely generated group on generators  $g_1, \dots, g_\ell$  satisfying

$$(g_i g_j)^{m_{ij}} = 1$$

where  $m_{ii} = 1$  and  $m_{ij} \geq 2$  for  $i \neq j$ . In [ČST18] the authors first showed that if  $G$  is a finite Coxeter group,  $\lambda$  is its left regular representation, and  $T = (g_1, \dots, g_\ell)$  are the generators as above, then  $\mathcal{Z}_1(T, \lambda)$  determines  $G$  up to isomorphism. Furthermore, if  $G$  is not of exceptional type (in the Coxeter diagram sense) and  $\rho$  is an arbitrary finite-dimensional representation of  $G$ , then  $\mathcal{Z}_1(T, \rho)$  determines  $\rho$ .

These theorems were presented during the Multivariable Spectral Theory and Representation Theory workshop at the Banff International Research Station in April 2019. Several problems about extending these results beyond Coxeter groups were posed by the speakers; among them were the following.

**Questions 1.1.** *Let  $G$  be a finite group,  $T$  a fixed generating set for  $G$ , and  $\rho_1, \rho_2$  irreducible complex representations of  $G$ .*

- (1) *Is  $\mathcal{Z}_1(T, \rho_1)$  a reduced and irreducible hypersurface?*
- (2) *If  $\mathcal{Z}_1(T, \rho_1) = \mathcal{Z}_1(T, \rho_2)$ , are  $\rho_1$  and  $\rho_2$  equivalent?*

As usual,  $\rho_1 : G \rightarrow \mathrm{GL}_{d_1}(\mathbb{C})$  and  $\rho_2 : G \rightarrow \mathrm{GL}_{d_2}(\mathbb{C})$  are equivalent if  $d_1 = d_2$  and  $\rho_2 = P \rho_1 P^{-1}$  for some  $P \in \mathrm{GL}_{d_1}(\mathbb{C})$ . A representation  $\rho_1$  is irreducible if its image does not admit a nontrivial common invariant subspace;

equivalently, it generates  $M_{d_1}(\mathbb{C})$  as a  $\mathbb{C}$ -algebra by Burnside's theorem [Isa76, Corollary 1.17]. The hypersurface  $\mathcal{X}_1(T, \rho_1)$  is reduced and irreducible (in the scheme-theoretic sense) if its defining determinant in (1.2) is an irreducible polynomial. The main result of this note is the following.

**Theorem 1.2.** *Questions 1.1(1) & (2) have negative answers in general.*

See Subsections 2.1 and 2.2 for concrete examples. On a more positive side, in Section 3 we show that representation theory aligns well with a quantization of the determinantal hypersurface, the free locus; see Theorem 3.1. Furthermore, Proposition 3.4 determines whether a free locus arises from a representation of a finite group, and Proposition 3.7 characterizes finite abelian groups from the perspective of determinantal hypersurfaces. We conclude this note with an open question.

## 2. Representations versus determinants

In this section we give negative answers to Questions 1.1. The representations were found with the help of the computer algebra system GAP and the online repository ATLAS of Finite Group Representations. Verifying equivalence and irreducibility of representations was sometimes done symbolically with the computing system Mathematica.

### 2.1. Irreducible representation with reducible determinant

The alternating group  $G = A_6$  admits a presentation

$$G = \langle g_1, g_2 \mid g_1^2, g_2^4, (g_1 g_2)^5, (g_1 g_2^2)^5 \rangle.$$

Let

$$A_1 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ -1 & -1 & -1 & -1 & -1 & -1 & -1 & -1 & -1 & -1 \end{pmatrix}$$

and

$$A_2 = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

Then  $\rho(g_1) = A_1$  and  $\rho(g_2) = A_2$  determines a faithful and irreducible representation  $\rho : G \rightarrow \mathrm{GL}_9(\mathbb{C})$ . Indeed, we can directly check that

$$A_1^2 = A_2^4 = (A_1 A_2)^5 = (A_1 A_2^2)^5 = I,$$

so  $\rho$  is a representation of  $G$ , and is moreover faithful since it is nontrivial and  $G$  is simple. Furthermore, all the possible products of  $A_1$  and  $A_2$  with at most 8 factors span the whole  $M_9(\mathbb{C})$ , so  $\rho$  is irreducible. However, we claim that the curve  $\mathcal{X}_1((g_1, g_2), \rho)$  in  $\mathbb{C}^2$  is not irreducible. We can compute the determinant of  $I + x_1 \rho(g_1) + x_2 \rho(g_2)$ ,

$$\det \begin{pmatrix} 1 + x_1 & x_2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & x_1 & x_2 & 0 & 0 & 0 & 0 & 0 \\ 0 & x_1 & 1 & 0 & x_2 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & x_1 & x_2 & 0 & 0 & 0 \\ 0 & 0 & 0 & x_1 & 1 & 0 & x_2 & 0 & 0 \\ x_2 & 0 & 0 & 0 & 0 & 1 & 0 & x_1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 + x_1 & 0 & x_2 \\ 0 & 0 & 0 & 0 & 0 & x_1 & 0 & 1 + x_2 & 0 \\ -x_1 & -x_1 & x_2 - x_1 & -x_1 & -x_1 & -x_1 & -x_1 & -x_1 & 1 - x_1 \end{pmatrix}$$

by cofactor expansion along the rows. The reader will have no difficulty verifying that  $\det(I + x_1 \rho(g_1) + x_2 \rho(g_2))$  equals

$$\begin{aligned} & 1 + x_1 - 4x_1^2 - 4x_1^3 + 6x_1^4 + 6x_1^5 - 4x_1^6 - 4x_1^7 + x_1^8 + x_1^9 + x_2 + 2x_1x_2 - 2x_1^2x_2 \\ & - 6x_1^3x_2 + 6x_1^5x_2 + 2x_1^6x_2 - 2x_1^7x_2 - x_1^8x_2 + x_1^2x_2^2 + x_1^3x_2^2 - 2x_1^4x_2^2 - 2x_1^5x_2^2 \\ & + x_1^6x_2^2 + x_1^7x_2^2 - x_1^2x_2^3 + 2x_1^4x_2^3 - x_1^6x_2^3 - 2x_2^4 + x_1^2x_2^4 - x_1^3x_2^4 + x_1^4x_2^4 + x_1^5x_2^4 \\ & - 2x_2^5 - 2x_1x_2^5 - x_1^2x_2^5 + x_1^4x_2^5 - x_1^2x_2^6 - x_1^3x_2^6 + x_1^2x_2^7 + x_2^8 - x_1x_2^8 + x_2^9 \end{aligned}$$

which is the product of the following two irreducible polynomials:

$$\begin{aligned} & 1 + 2x_1 - 2x_1^3 - x_1^4 + x_1x_2 + 2x_1^2x_2 + x_1^3x_2 - x_1x_2^2 - x_1^2x_2^2 + x_1x_2^3 - x_2^4, \\ & 1 - x_1 - 2x_1^2 + 2x_1^3 + x_1^4 - x_1^5 + x_2 - x_1x_2 - x_1^2x_2 + x_1^3x_2 - x_2^4 - x_2^5. \end{aligned}$$

Some of the subsequent examples are presented in a more terse way to maintain the focus on their intent.

Note that the above irreducible representation of  $A_6$  has dimension 9, which is not the minimum among nontrivial complex representations of  $A_6$ ; namely,  $A_6$  admits a representation  $\sigma$  of minimal dimension 5, and  $\mathcal{X}_1((g_1, g_2), \sigma)$  is irreducible. One might thus be tempted to suggest that for a group  $G$  generated by a finite set  $T$  and its (irreducible) representation  $\sigma$  of minimal dimension,  $\mathcal{X}_1(T, \sigma)$  is irreducible. However, even this weaker conjecture fails. The counterexample is given by the Janko group  $J_2$ ,

$$J_2 = \langle g_1, g_2 \mid g_1^2, g_2^3, (g_1g_2)^7, (g_1g_2g_1^{-1}g_2^{-1})^{12}, (g_1g_2(g_1g_2g_1g_2^{-1})^2)^6 \rangle.$$

This sporadic simple group of order 604800 admits two non-isomorphic complex representations  $\sigma_1, \sigma_2$  of minimal dimension 14, courtesy of ATLAS of Finite Group Representations. As in the previous example (albeit with slightly

longer calculations), one can explicitly check that the curve  $\mathcal{Z}_1((g_1, g_2), \sigma_1) = \mathcal{Z}_1((g_1, g_2), \sigma_2)$  has two irreducible components.

### 2.2. Non-equivalent representations with the same determinant

The classical group  $G = \mathrm{GL}_2(\mathbb{Z}/3\mathbb{Z})$  admits the presentation

$$G = \langle g_1, g_2, g_3 \mid g_1^2, (g_1 g_2^{-1})^2, (g_1 g_3^{-1})^2, g_2^2 g_3 g_2^{-1} g_3, g_2 g_3^2 g_2 g_3^{-1} \rangle.$$

Let  $A_1, A_2, A_3$  be the matrices

$$\begin{pmatrix} -\frac{1}{\sqrt{2}} & -\frac{1}{2} - \frac{i}{2} \\ -\frac{1}{2} + \frac{i}{2} & \frac{1}{\sqrt{2}} \end{pmatrix}, \quad \begin{pmatrix} \frac{1}{2} + \frac{i}{2} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & \frac{1}{2} - \frac{i}{2} \end{pmatrix}, \quad \begin{pmatrix} \frac{1}{2} - \frac{i}{2} & \frac{i}{\sqrt{2}} \\ \frac{i}{\sqrt{2}} & \frac{1}{2} + \frac{i}{2} \end{pmatrix}.$$

There are faithful irreducible unitary representations  $\rho_+, \rho_- : G \rightarrow \mathrm{GL}_2(\mathbb{C})$  given by

$$\rho_{\pm}(g_1) = \pm A_1, \quad \rho_{\pm}(g_2) = A_2, \quad \rho_{\pm}(g_3) = A_3.$$

It is easy to check that  $\rho_+$  and  $\rho_-$  are not equivalent. On the other hand,

$$\mathcal{Z}_1((g_1, g_2, g_3), \rho_{\pm}) = \{(\xi_1, \xi_2, \xi_3) : 1 - \xi_1^2 + \xi_2 + \xi_2^2 + \xi_3 + \xi_3^2 = 0\}.$$

## 3. Free locus perspective

In this section we will see how representations of a finitely generated group are determined by a noncommutative relaxation of (1.2). To  $A \in M_d(\mathbb{C})^\ell$  we associate the monic matrix pencil  $L_A = I + A_1 x_1 + \cdots + A_\ell x_\ell$  of size  $d$  in freely noncommuting variables  $x = (x_1, \dots, x_\ell)$ . Thus  $L$  is an affine matrix over the free algebra  $\mathbb{C}\langle x \rangle$ . At a matrix point  $X \in M_n(\mathbb{C})^\ell$  it evaluates as

$$L_A(X) = I_{dn} + A_1 \otimes X_1 + \cdots + A_\ell \otimes X_\ell \in M_{dn}(\mathbb{C}).$$

The **free locus** [KV17] of  $L_A$  is the disjoint union of determinantal hypersurfaces

$$\mathcal{Z}(L_A) = \bigsqcup_{n \in \mathbb{N}} \mathcal{Z}_n(L_A), \quad \mathcal{Z}_n(L_A) = \{X \in M_n(\mathbb{C})^\ell : \det L_A(X) = 0\}.$$

Given a group  $G$  generated by  $T = (g_1, \dots, g_n)$  and a complex representation  $\rho : G \rightarrow \mathrm{GL}_d(\mathbb{C})$ , we write

$$\mathcal{Z}(T, \rho) = \mathcal{Z}(L_{\rho(g_1), \dots, \rho(g_\ell)}). \tag{3.1}$$

By the definition of the free locus we see that (3.1) is indeed a quantization of (1.2). The existing results on free loci [KV17, HKV18] readily apply to group representations.

**Theorem 3.1.** *For  $i = 1, 2$  let  $G_i$  be a group generated by a finite sequence  $T_i$  and let  $\rho_i$  be a complex representation of  $G_i$ . Assume  $|T_1| = |T_2|$ .*

- (1) *If  $\rho_i$  is irreducible, then there exists  $n_0 \in \mathbb{N}$  such that  $\mathcal{Z}_n(T_i, \rho_i)$  is a reduced and irreducible hypersurface for all  $n \geq n_0$ .*
- (2) *If  $\rho_1$  and  $\rho_2$  are irreducible, then  $\mathcal{Z}(T_1, \rho_1) = \mathcal{Z}(T_2, \rho_2)$  if and only if  $G_1 / \ker \rho_1 \cong G_2 / \ker \rho_2$  and  $\rho_1, \rho_2$  are equivalent.*

- (3) For  $i = 1, 2$  assume that  $G_i$  is finite and  $\rho_i$  is a faithful representation. Then  $\mathcal{Z}(T_1, \rho_1) = \mathcal{Z}(T_2, \rho_2)$  if and only if  $G_1 \cong G_2$  via an isomorphism mapping  $T_1$  to  $T_2$ .

*Proof.* (1) A consequence of [HKV18, Theorem 3.4].

(2) A consequence of [KV17, Theorem 3.11].

(3) Let  $\mathcal{T}_i$  be the  $\mathbb{C}$ -algebra generated by  $T_i$ . Since  $G_i$  is finite, its group algebra  $\mathbb{C}G_i$  is semisimple by Maschke's theorem [Isa76, Theorem 1.9]. Since  $\mathcal{T}_i$  is a quotient of  $\mathbb{C}G_i$ , it is also semisimple. Then  $\mathcal{Z}(T_1, \rho_1) = \mathcal{Z}(T_2, \rho_2)$  if and only if  $T_1 \mapsto T_2$  induces an algebra isomorphism  $\mathcal{T}_1 \rightarrow \mathcal{T}_2$  by [KV17, Corollary 3.8]. This isomorphism then restricts to the group isomorphism  $G_1 \rightarrow G_2$ .  $\square$

*Remark 3.2.* There is a deterministic bound on  $n_0$  in Theorem 3.1(1) that is exponential in  $|T_1|$  and the dimension of  $\rho_1$  by [HKV18, Remark 3.5] (the bound is likely not optimal). Similarly, to verify  $\mathcal{Z}(T_1, \rho_1) = \mathcal{Z}(T_2, \rho_2)$  of Theorem 3.1(2,3), it suffices to check  $\mathcal{Z}_n(T_1, \rho_1) = \mathcal{Z}_n(T_2, \rho_2)$  for a fixed large enough  $n$ , exponential in  $|T_i|$  and the dimension of  $\rho_i$  by [KV17, Remark 3.7].

Free loci are defined for monic pencils with arbitrary matrix coefficients; we now describe how the geometry of the free locus  $\mathcal{Z}(L_A)$  detects whether the coefficients  $A_1, \dots, A_\ell$  generate a finite group. See also [DFO'B13] for an efficient algorithm that determines finiteness of a finitely generated linear group.

**Definition 3.3.** Let  $\ell, n \in \mathbb{N}$ . Let  $C \in \mathrm{GL}_n(\mathbb{Z})$  be the permutation matrix corresponding to the cycle  $(1\ 2\ \dots\ n)$ . If  $\{1, \dots, n\} = S_1 \sqcup \dots \sqcup S_\ell$  and  $P_j$  is the orthogonal projection onto  $\mathrm{span}\{e_k : k \in S_j\}$ , then the matrix point

$$X = (P_1 C, \dots, P_\ell C) \in \mathrm{M}_n(\mathbb{Z})^\ell$$

is called a **cycle partition**. For given  $\ell, n$  we thus have  $\ell^n$  cycle partitions.

Let  $\mu_\infty \subset \mathbb{C} \setminus \{0\}$  be the group of all roots of unity. The next proposition shows that if  $A_1, \dots, A_\ell$  generate a finite group, then  $\mathcal{Z}(L_A)$  intersects complex lines through cycle partitions only in points from  $\mu_\infty$ .

**Proposition 3.4.** *Let  $A \in \mathrm{M}_d(\mathbb{C})^\ell$ . Then  $A_1, \dots, A_\ell$  generate a finite group if and only if the following hold:*

- (i) *there is a positive definite  $P \in \mathrm{M}_d(\mathbb{C})$  such that  $A_j^* P A_j = P$  for all  $j$ ;*
- (ii) *for every cycle partition  $X$  and  $t \in \mathbb{C}$ ,*

$$tX \in \mathcal{Z}(L_A) \implies t \in \mu_\infty.$$

*Proof.* ( $\Leftarrow$ ) Every  $A_j$  is invertible by (i). Let  $G$  be a group generated by  $A_1, \dots, A_\ell$ . Also by (i),  $G$  is a subgroup of the unitary group in  $\mathrm{GL}_d(\mathbb{C})$  with respect to the inner product  $\langle u, v \rangle = u^* P v$ . Hence every element of  $G$  is diagonalizable. By [Weh73, Corollary 4.9], a finitely generated subgroup of  $\mathrm{GL}_d(\mathbb{C})$  is finite if and only if it is periodic (or torsion; i.e., every element has finite order). Since a diagonalizable matrix has a finite order if and only

if all its eigenvalues lie in  $\mu_\infty$ , it suffices to verify that eigenvalues of every element of  $G$  lie in  $\mu_\infty$ .

To  $(i_1, \dots, i_n) \in \{1, \dots, \ell\}^n$  we associate the cycle partition  $X \in M_n(\mathbb{Z})^\ell$  by choosing  $S_j = \{e_k : i_k = j\}$ . We claim that  $tX \in \mathcal{L}(L_A)$  if and only if  $(-t)^n$  is an eigenvalue of  $A_{i_1} \cdots A_{i_n}$ . Indeed, using Schur complements it is easy to check that

$$\begin{aligned} \det(I - (-1)^n t^n A_{i_1} \cdots A_{i_n}) &= \det \begin{pmatrix} I & & & tA_{i_1} \\ (-1)^n t^{n-1} A_{i_2} \cdots A_{i_n} & & & I \end{pmatrix} \\ &= \det \begin{pmatrix} I & & tA_{i_1} & 0 \\ 0 & & I & tA_{i_2} \\ -(-1)^n t^{n-2} A_{i_3} \cdots A_{i_n} & & 0 & I \end{pmatrix} \\ &= \dots \\ &= \det \begin{pmatrix} I & tA_{i_1} & & & \\ & \ddots & \ddots & & \\ & & I & tA_{i_{n-1}} & \\ tA_{i_n} & & & I & \end{pmatrix} \\ &= \det L_A(tX). \end{aligned}$$

Thus the matrix  $A_{i_1} \cdots A_{i_n}$  has finite order if and only if  $tX \in \mathcal{L}(L_A)$  implies  $t \in \mu_\infty$ , which holds by (ii).

( $\Rightarrow$ ) If  $A_1, \dots, A_\ell$  generate a finite group  $G$ , then  $\mathbb{C}^d$  admits a  $G$ -invariant inner product

$$\langle u, v \rangle = \sum_{g \in G} (gu)^*(gv).$$

If  $P$  is the positive definite matrix satisfying  $\langle u, v \rangle = u^* P v$ , then (i) holds. Furthermore, the proof of (ii) is already given in the previous paragraph.  $\square$

*Remark 3.5.* If additional information about  $A_1, \dots, A_\ell$  is given, say that their entries generate a number field (finite extension of  $\mathbb{Q}$ ), then the size of the cycle partitions, which have to be tested in Proposition 3.4, can be bounded using Schur's theorem on orders of finite matrix groups [Isa76, Theorem 14.19].

*Remark 3.6.* Let  $p \in \mathbb{N}$  be prime. If  $\mu_\infty$  in Proposition 3.4 is replaced by the group of power-of- $p$  roots of unity, one obtains a free locus characterization of matrix tuples that generate a finite  $p$ -group.

We also show how the free locus certifies whether its defining coefficients generate a finite abelian group. The degree of an affine variety of codimension  $m$  is the number of intersection points of the variety with  $m$  hyperplanes in general position; in the case of a hypersurface, it is simply the degree of its square-free defining polynomial.

**Proposition 3.7.** *Let  $G$  be a finite group generated by  $A_1, \dots, A_\ell \in M_d(\mathbb{C})$ . Then  $G$  is abelian if and only if the irreducible components of  $\mathcal{L}_n(L_A)$  have degree  $n$  for all  $n \in \mathbb{N}$ .*

*Proof.* Let  $\mathcal{A}$  be the  $\mathbb{C}$ -algebra generated by  $A_1, \dots, A_\ell$ . As in the proof of Theorem 3.1(3) we see that  $\mathcal{A}$  is semisimple. After a basis change (which does not affect the structure of  $G$  or  $\mathcal{Z}(L_A)$ ) we can thus assume that

$$A_j = A_j^{(1)} \oplus \dots \oplus A_j^{(s)}$$

where  $A_1^{(k)}, \dots, A_\ell^{(k)} \in M_{d_k}(\mathbb{C})$  determine an irreducible representation of  $G$  for every  $k = 1, \dots, s$ . For  $X \in M_n(\mathbb{C})^d$  let us view  $\det L_{A^{(k)}}(X)$  as a polynomial in the entries of  $X$ . If  $d_k = 1$ , then  $\det L_{A^{(k)}}(X)$  is up to an affine change of coordinates equal to the determinant of a generic  $n \times n$  matrix, and hence an irreducible polynomial of degree  $n$ . On the other hand, if  $d_k > 1$ , then  $\det L_{A^{(k)}}(X)$  is a polynomial of degree  $d_k n > n$  for all  $n$ , and irreducible for all large enough  $n$  by [HKV18, Theorem 3.4]. Since  $G$  is abelian if and only if  $d_1 = \dots = d_s = 1$ , and

$$\mathcal{Z}_n(L_A) = \mathcal{Z}_n(L_A^{(1)}) \cup \dots \cup \mathcal{Z}_n(L_A^{(s)}),$$

it follows that  $G$  is abelian if and only if the irreducible components of  $\mathcal{Z}_n(L_A)$  are hypersurfaces of degree  $n$ .  $\square$

*Remark 3.8.* If  $\ell = 2$  and  $A_1, A_2$  are hermitian, then  $\mathcal{Z}_1(L_A)$  alone determines whether  $G$  is abelian, cf. [MT52].

The last two propositions offer some directions for future research. Theorem 3.1 implies that the linear group  $G$  generated by a tuple  $A$  is determined by  $\mathcal{Z}(L_A)$ . It would be interesting to know which properties of  $G$  can be deduced from the geometry of  $\mathcal{Z}(L_A)$ . For example, intersections of  $\mathcal{Z}(L_A)$  with certain lines and hyperplanes determine whether  $G$  is finite or abelian. An open problem is how to decide whether a finite group  $G$  is nilpotent/solvable/simple (or any other group-theoretic property) by considering the geometry of the hypersurfaces  $\mathcal{Z}_n(L_A)$ .

### Acknowledgment

The authors thank Banff International Research Station for the hospitality during the Multivariable Spectral Theory and Representation Theory workshop, and participants for sharing their ideas.

### References

- [Bea00] A. Beauville: *Determinantal hypersurfaces*, Michigan Math. J. 48 (2000) 39–64.
- [Brä11] P. Brändén: *Obstructions to determinantal representability*, Adv. Math. 226 (2011) 1202–1212.
- [CY13] P. Cade, R. Yang: *Projective spectrum and cyclic cohomology*, J. Funct. Anal. 265 (2013) 1916–1933.
- [CSZ16] I. Chagouel, M. I. Stessin, K. Zhu: *Geometric spectral theory for compact operators*, Trans. Amer. Math. Soc. 368 (2016) 1559–1582.
- [ČST18] Ž. Čučković, M. I. Stessin, A. B. Tchernev: *Determinantal hypersurfaces and representations of Coxeter groups*, preprint arXiv:1810.12893.



- [dCEP18] C. de Concini, D. Eisenbud, C. Procesi: *Young diagrams and determinantal varieties*, Invent. Math. 56 (1980) 129–165.
- [Ded68] R. Dedekind: *Gesammelte mathematische Werke*, Band II, Chelsea Publishing Co., New York, 1968.
- [DFO'B13] A. S. Detinko, D. L. Flannery, E. A. O'Brien: *Recognizing finite matrix groups over infinite fields*, J. Symbolic Comput. 50 (2013) 100–109.
- [Dic02] L. E. Dickson: *An elementary exposition of Frobenius's theory of group characters and group-determinants*, Ann. of Math. 4 (1902) 25–49.
- [Dic21] L. E. Dickson: *Determination of all general homogeneous polynomials expressible as determinants with linear elements*, Trans. Amer. Math. Soc. 22 (1921) 167–179.
- [Dol12] I. Dolgachev: *Classical algebraic geometry: a modern view*, Cambridge University Press, Cambridge, 2012.
- [FS91] E. Formanek, D. Sibley: *The group determinant determines the group*, Proc. Amer. Math. Soc. 112 (1991) 649–656.
- [Fro68] F. G. Frobenius: *Gesammelte Abhandlungen*, Bände I–III, Springer-Verlag, Berlin-New York, 1968.
- [GY17] R. I. Grigorchuk, R. Yang: *Joint spectrum and the infinite dihedral group*, Proc. Steklov Inst. Math. 297 (2017) 145–178.
- [HKV18] J.W. Helton, I. Klep, J. Volčič: *Geometry of free loci and factorization of noncommutative polynomials*, Adv. Math. 331 (2018) 589–626.
- [HV07] J. W. Helton, V. Vinnikov: *Linear matrix inequality representation of sets*, Comm. Pure Appl. Math. 60 (2007) 654–674.
- [Isa76] I. M. Isaacs: *Character theory of finite groups*, Pure and Applied Mathematics 69, Academic Press, New York–London, 1976.
- [KV12] D. Kerner, V. Vinnikov: *Determinantal representations of singular hypersurfaces in  $\mathbb{P}^n$* , Adv. Math. 231 (2012) 1619–1654.
- [KV17] I. Klep, J. Volčič: *Free loci of matrix pencils and domains of noncommutative rational functions*, Comment. Math. Helv. 92 (2017) 105–130.
- [MT52] T. S. Motzkin, O. Taussky: *Pairs of matrices with property L*, Trans. Amer. Math. Soc. 73 (1952) 108–114.
- [ST15] M. I. Stessin, A. B. Tchernev: *Spectral algebraic curves and decomposable operator tuples*, J. Operator Theory 82 (2019) 75–113.
- [Tay70] J. L. Taylor: *A joint spectrum for several commuting operators*, J. Funct. Anal. 6 (1970) 172–191.
- [Vol19] J. Volčič: *Stable noncommutative polynomials and their determinantal representations*, SIAM J. Appl. Algebra Geometry 3 (2019) 152–171.
- [Wag11] D. G. Wagner: *Multivariate stable polynomials: theory and applications*, Bull. Amer. Math. Soc. (N.S.) 48 (2011) 53–84.
- [Weh73] B. A. F. Wehrfritz: *Infinite linear groups*, Ergebnisse der Mathematik und ihrer Grenzgebiete 76, Springer-Verlag, New York-Heidelberg, 1973.
- [Yan09] R. Yang: *Projective spectrum in Banach algebras*, J. Topol. Anal. 1 (2009) 289–306.

Igor Klep  
Department of Mathematics  
Faculty of Mathematics and Physics  
University of Ljubljana  
Slovenia  
e-mail: [igor.klep@fmf.uni-lj.si](mailto:igor.klep@fmf.uni-lj.si)

Jurij Volčič  
Department of Mathematics  
Texas A&M University  
USA  
e-mail: [volcic@math.tamu.edu](mailto:volcic@math.tamu.edu)