

# INFEASIBILITY CERTIFICATES FOR LINEAR MATRIX INEQUALITIES

IGOR KLEP AND MARKUS SCHWEIGHOFER

ABSTRACT. Farkas' lemma is a fundamental result from *linear programming* providing *linear* certificates for infeasibility of systems of linear inequalities. In *semidefinite programming*, such linear certificates only exist for *strongly* infeasible linear matrix inequalities. We provide *nonlinear algebraic* certificates for *all* infeasible linear matrix inequalities in the spirit of real algebraic geometry. More precisely, we show that a linear matrix inequality  $L(x) \succeq 0$  is infeasible if and only if  $-1$  lies in the quadratic module associated to  $L$ . We prove exponential degree bounds for the corresponding algebraic certificate. In order to get a polynomial size certificate, we use a more involved algebraic certificate motivated by the real radical and Prestel's theory of semiorderings. Completely different methods, namely complete positivity from operator algebras, are employed to consider linear matrix inequality domination.

A *linear matrix inequality* (LMI) is a condition of the form

$$L(x) = A_0 + \sum_{i=1}^n x_i A_i \succeq 0 \quad (x \in \mathbb{R}^n)$$

where the  $A_i$  are symmetric matrices of the same size and one is interested in the solutions  $x \in \mathbb{R}^n$  making  $L(x) \succeq 0$ , i.e., making  $L(x)$  into a positive semidefinite matrix. The solution set to such an inequality is a closed convex semialgebraic subset of  $\mathbb{R}^n$  called a *spectrahedron*. Optimization of linear objective functions over spectrahedra is called *semidefinite programming* (SDP) [BV, To, WSV]. In this article, we are mainly concerned with the important SDP feasibility problem: When is an LMI *feasible*; i.e., when is there an  $x \in \mathbb{R}^n$  such that  $L(x) \succeq 0$ ?

Note that a diagonal LMI (all  $A_i$  diagonal matrices) is just a (finite) system of (non-strict) linear inequalities. The solution set of such a linear system is a (closed convex) polyhedron. Optimization of linear objective functions over polyhedra is called *linear programming* (LP). The ellipsoid method developed by Shor, Yudin, Nemirovskii and Khachiyan showed at the end of the 1970s for the first time that the LP feasibility problem (and actually the problem of solving LPs) can be solved in (deterministically) polynomial time (in the bit model of computation assuming rational coefficients) [Sr, Chapter 13]. Another breakthrough came in the 1980s with the introduction of the more practical interior point methods by Karmarkar and their theoretical underpinning by Nesterov and Nemirovskii [NN, Ne].

The motivation to replace the prefix “poly” by “spectra” is to replace “many” values of linear polynomials (the diagonal values of  $L(x)$ ) by the “spectrum” of

---

*Date:* 25 August 2011.

*2010 Mathematics Subject Classification.* Primary 13J30, 15A22, 46L07, 90C22; Secondary 14P10, 15A48.

*Key words and phrases.* linear matrix inequality, LMI, spectrahedron, semidefinite program, quadratic module, infeasibility, duality, complete positivity, Farkas' lemma.

This research was supported through the programme “Research in Pairs” (RiP) by the Mathematisches Forschungsinstitut Oberwolfach in 2010. The first author was also supported by the Slovenian Research Agency under Project no. J1-3608 and Program no. P1-0222. Part of the research was done while the first author held a visiting professorship at the Universität Konstanz in 2011.

$L(x)$  (i.e., the set of its eigenvalues, all of which are real since  $L(x)$  is symmetric). The advantage of LMIs over systems of linear inequalities (or of spectrahedra over polyhedra, and SDP over LP, respectively) is a considerable gain of expressiveness which makes LMIs an important tool in many areas of applied and pure mathematics. Many problems in control theory, system identification and signal processing can be formulated using LMIs [BEFB, Par, SIG]. Combinatorial optimization problems can often be modeled or approximated by SDPs [Go]. LMIs also find application in real algebraic geometry for finding sums of squares decompositions of polynomials [Las, Ma]. Strongly related to this, there is a hierarchy of SDP approximations to polynomial optimization problems [Lau] consisting of the so-called Lasserre moment relaxations. This hierarchy and related methods recast the field of polynomial optimization (where the word “polynomial” stands for polynomial objective and polynomial constraints). In this article, rather than trying to solve polynomial optimization problems by using SDPs, we borrow ideas and techniques from real algebraic geometry and polynomial optimization in order to get new results in the theory of semidefinite programming.

The price to pay for the increased expressivity of SDPs is that they enjoy some less good properties. The complexity of solving general SDPs is a very subtle issue which is often downplayed. For applications in combinatorial optimization, it follows typically from the general theory of the ellipsoid method [Sr] or interior point methods [NN] that the translation into SDPs yields a polynomial time algorithm (see for instance [dK, Section 1.9] for exact statements). However the complexity status of the LMI feasibility problem (the problem of deciding whether a given LMI with rational coefficients has a solution) is largely unknown. What is known is essentially only that (in the bit model) LMI feasibility lies either in  $\text{NP} \cap \text{co-NP}$  or outside  $\text{NP} \cup \text{co-NP}$ . This will also follow from our work below, but has been already proved by Ramana [Ra] in 1997. The standard (Lagrange-Slater) dual of a semidefinite program works well when the feasible set is full-dimensional (e.g. if there is  $x \in \mathbb{R}^n$  with  $L(x) \succ 0$ ). However, in general strong duality can fail badly which is a serious problem since there is no easy way of reducing to the full-dimensional case (cf. the LMI feasibility problem mentioned above). Even the corresponding version of Farkas’ lemma fails for SDP.

Ramana’s as well as our proof relies on an extension of the standard (Lagrange-Slater) dual of an SDP which can be produced in polynomial time from the primal and for which strong duality (more precisely zero gap and dual attainment) always holds. Ramana’s extension is an encoding of the standard dual of a regularized primal SDP in the sense of Borwein and Wolkowicz [BW]. Our dual, although having some superficial similarity to Ramana’s, relies on completely different ideas, namely sums of squares certificates of nonnegativity. The ideas for this *sums of squares dual* come from real algebraic geometry, more precisely from sums of squares representations and the Real Nullstellensatz [Ma, PD, Sc]. We believe that this new connection will lead to further insights in the future.

The paper is organized as follows: We fix terminology and notation in Section 1. Our first line of results is given in Section 2, where we give an algebraic characterization of infeasible LMIs (see Theorem 2.2.5 and Corollary 2.3.3) involving the quadratic module associated to an LMI. Our characterization allows us to construct a new LMI whose feasibility is equivalent to the infeasibility of the original LMI. This new LMI is canonical from the viewpoint of positive polynomials. However its size is exponential in the size of the primal, so in Section 3 we use real algebraic geometry to construct a polynomial size LMI whose feasibility is equivalent to the infeasibility of the original LMI, cf. Theorem 3.5.2. At the same time Theorem

3.5.2 gives a new type of a linear Positivstellensatz characterizing linear polynomials nonnegative on a spectrahedron. The article concludes with Section 4, where we revisit the [HKM] noncommutative (matricial) relaxation of an LMI and investigate how our duality theory of SDP pertains to (completely) positive maps.

## 1. NOTATION AND TERMINOLOGY

We write  $\mathbb{N} := \{1, 2, \dots\}$ ,  $\mathbb{Q}$ , and  $\mathbb{R}$  for the sets of natural, rational, and real numbers, respectively. For any matrix  $A$ , we denote by  $A^*$  its transpose.

**1.1. Sums of squares.** Let  $R$  be a (commutative unital) ring. Then  $SR^{m \times m} := \{A \in R^{m \times m} \mid A = A^*\}$  denotes the set of all *symmetric*  $m \times m$  matrices. Examples of these include *hermitian squares*, i.e., elements of the form  $A^*A$  for some  $A \in R^{m \times m}$ .

Recall that a matrix  $A \in \mathbb{R}^{m \times m}$  is called *positive semidefinite* (*positive definite*) if it is symmetric and  $v^*Av \geq 0$  for all (column) vectors  $v \in \mathbb{R}^m$ ,  $A$  is *positive definite* if it is positive semidefinite and invertible. For real matrices  $A$  and  $B$  of the same size, we write  $A \preceq B$  (respectively  $A \prec B$ ) to express that  $B - A$  is positive semidefinite (respectively positive definite). We denote by  $S\mathbb{R}_{\succeq 0}^{m \times m}$  and  $S\mathbb{R}_{\succ 0}^{m \times m}$  the convex cone of all positive semidefinite and positive definite matrices of size  $m$ , respectively.

Let  $\underline{X} = (X_1, \dots, X_n)$  be an  $n$ -tuple of commuting variables and  $\mathbb{R}[\underline{X}]$  the polynomial ring. With  $\mathbb{R}[\underline{X}]_k$  we denote the vector space of all polynomials of degree  $\leq k$ , and  $\sum \mathbb{R}[\underline{X}]^2$  is the convex cone of all sums of squares (*sos-polynomials*), i.e.,

$$\sum \mathbb{R}[\underline{X}]^2 = \left\{ \sum_{i=1}^r p_i^2 \mid r \in \mathbb{N}, p_i \in \mathbb{R}[\underline{X}] \right\}.$$

A (real) *matrix polynomial* is a matrix whose entries are polynomials from  $\mathbb{R}[\underline{X}]$ . It is *linear* or *quadratic* if its entries are from  $\mathbb{R}[\underline{X}]_1$  or  $\mathbb{R}[\underline{X}]_2$ , respectively. A matrix polynomial is called *symmetric* if it coincides with its transpose. An example of symmetric matrix polynomials that are of special interest to us are sums of hermitian squares in  $\mathbb{R}[\underline{X}]^{m \times m}$ . They are called *sos-matrices*. More explicitly,  $S \in \mathbb{R}[\underline{X}]^{m \times m}$  is an *sos-matrix* if the following equivalent conditions hold:

- (i)  $S = P^*P$  for some  $s \in \mathbb{N}$  and some  $P \in \mathbb{R}[\underline{X}]^{s \times m}$ ;
- (ii)  $S = \sum_{i=1}^r Q_i^*Q_i$  for some  $r \in \mathbb{N}$  and  $Q_i \in \mathbb{R}[\underline{X}]^{m \times m}$ ;
- (iii)  $S = \sum_{i=1}^s v_i v_i^*$  for some  $s \in \mathbb{N}$  and  $v_i \in \mathbb{R}[\underline{X}]^m$ .

Note that an *sos-matrix*  $S \in \mathbb{R}[\underline{X}]^{m \times m}$  is positive semidefinite on  $\mathbb{R}^n$  but not vice-versa, since e.g. a polynomial nonnegative on  $\mathbb{R}^n$  is not necessarily a sum of squares of polynomials [Ma, PD].

For a comprehensive treatment of the theory of matrix polynomials we refer the reader to the book [GLR] and the references therein.

**1.2. Linear pencils, spectrahedra, and quadratic modules.** We use the term *linear pencil* as a synonym and abbreviation for symmetric linear matrix polynomial.

Let  $R$  be a (commutative unital) ring. We recall that in real algebraic geometry a subset  $M \subseteq R$  is called a *quadratic module* in  $R$  if it contains 1 and is closed under addition and multiplication with squares, i.e.,

$$1 \in M, \quad M + M \subseteq M \quad \text{and} \quad a^2M \subseteq M \quad \text{for all } a \in R,$$

see for example [Ma]. A quadratic module  $M \subseteq R$  is called *proper* if  $-1 \notin M$ . Note that an improper quadratic module  $M$  in a ring  $R$  with  $\frac{1}{2} \in R$  equals  $R$  by the identity

$$4a = (a + 1)^2 - (a - 1)^2 \quad \text{for all } a \in R. \quad (1)$$

An LMI  $L(x) \succeq 0$  can be seen as the infinite family of simultaneous linear inequalities  $u^*L(x)u \geq 0$  ( $u \in \mathbb{R}^m$ ). In optimization, when dealing with families of linear inequalities, one often considers the convex cone generated by them (cf.  $C_L$  in the definition below). Real algebraic geometry handles *arbitrary* polynomial inequalities and uses the multiplicative structure of the polynomial ring. Thence one considers more special types of convex cones like in our case quadratic modules ( $M_L$  from the next definition). One of the aims of this article is to show that it is advantageous to consider quadratic modules for the study of LMIs. Since quadratic modules are infinite-dimensional convex cones, we will later on also consider certain finite-dimensional truncations of them, see Subsection 2.3.

**Definition 1.2.1.** Let  $L$  be a linear pencil of size  $m$  in the variables  $\underline{X}$ . We introduce

$$\begin{aligned} S_L &:= \{x \in \mathbb{R}^n \mid L(x) \succeq 0\} \subseteq \mathbb{R}^n, \\ C_L &:= \left\{ c + \sum_i u_i^* L u_i \mid c \in \mathbb{R}_{\geq 0}, u_i \in \mathbb{R}^m \right\} \\ &= \left\{ c + \text{tr}(LS) \mid c \in \mathbb{R}_{\geq 0}, S \in S\mathbb{R}_{\geq 0}^{m \times m} \right\} \subseteq \mathbb{R}[\underline{X}]_1, \\ M_L &:= \left\{ s + \sum_i v_i^* L v_i \mid s \in \sum \mathbb{R}[\underline{X}]^2, v_i \in \mathbb{R}[\underline{X}]^m \right\} \\ &= \left\{ s + \text{tr}(LS) \mid s \in \mathbb{R}[\underline{X}] \text{ sos-polynomial}, S \in \mathbb{R}[\underline{X}]^{m \times m} \text{ sos-matrix} \right\} \subseteq \mathbb{R}[\underline{X}], \end{aligned}$$

and call them the *spectrahedron* (or LMI set), the *convex cone*, and the *quadratic module* associated to the linear pencil  $L$ , respectively. Call the *linear matrix inequality* (LMI)

$$L(x) \succeq 0 \quad (x \in \mathbb{R}^n),$$

or simply  $L$ , *infeasible* if  $S_L = \emptyset$ . In this case, call it *strongly infeasible* if

$$\text{dist}(\{L(x) \mid x \in \mathbb{R}^n\}, S\mathbb{R}_{\geq 0}^{m \times m}) > 0,$$

and *weakly infeasible* otherwise. Moreover, call  $L$  *feasible* if  $S_L \neq \emptyset$ . We say  $L$  is *strongly feasible* if there is an  $x \in \mathbb{R}^n$  such that  $L(x) \succ 0$  and *weakly feasible* otherwise.

Note that for any linear pencil  $L$ , each element of  $M_L$  is a polynomial nonnegative on the spectrahedron  $S_L$ . In general  $M_L$  does not contain all polynomials nonnegative on  $S_L$ , e.g. when  $L$  is diagonal and  $\dim S_L \geq 3$  [Sc]. For diagonal  $L$ , the quadratic module  $M_L$  (and actually the convex cone  $C_L$ ) contains however all *linear* polynomials nonnegative on the polyhedron  $S_L$  by Farkas' lemma. For non-diagonal linear pencils  $L$  even this can fail, see Example 3.6.1 below. To certify nonnegativity of a linear polynomial on a spectrahedron, we therefore employ more involved algebraic certificates (motivated by the real radical), see Theorem 3.5.2.

## 2. SPECTRAHEDRA AND QUADRATIC MODULES

In this section we establish algebraic certificates for infeasibility and boundedness of a spectrahedron. They involve classical Positivstellensatz-like certificates expressing  $-1$  as a weighted sum of squares.

A quadratic module  $M \subseteq \mathbb{R}[\underline{X}]$  is said to be *archimedean* if

$$\forall f \in \mathbb{R}[\underline{X}] \exists N \in \mathbb{N} : N - f^2 \in M.$$

Equivalently, there is an  $N \in \mathbb{N}$  with  $N \pm X_i \in M$  for  $i = 1, \dots, n$ ; see [Ma, Corollary 5.2.4].

**2.1. Quadratic modules describing non-empty bounded spectrahedra.** Obviously, if  $M_L$  is archimedean for a linear pencil  $L$ , then  $S_L$  is bounded. In [HKM] complete positivity was used to deduce that for *strictly feasible* linear pencils the converse holds. Subsequently, a certain generalization of this result for *projections* of spectrahedra has been proved by other techniques in [GN]. In this section we will establish the result for arbitrary bounded  $S_L$ . We begin with the relatively easy case of non-empty  $S_L$  (possibly with empty interior).

**Lemma 2.1.1.** *Let  $L$  be a linear pencil with  $S_L \neq \emptyset$ . Then*

$$S_L \text{ is bounded} \iff M_L \text{ is archimedean.}$$

*Proof.* The direction ( $\Leftarrow$ ) is obvious as remarked above. Let us consider the converse. We first establish the existence of finitely many linear polynomials in  $C_L$  certifying the boundedness of  $S_L$ .

There is a ball  $B \subseteq \mathbb{R}^n$  with  $S_L \subseteq B$  and  $S_L \cap \partial B = \emptyset$ . For every  $x \in \partial B$  there is a vector  $u \in \mathbb{R}^n$  with

$$u^* L(x) u < 0. \quad (2)$$

By continuity, (2) holds for all  $x$  in a neighborhood  $U_x$  of  $x$ . From  $\{U_x \mid x \in \partial B\}$  we extract by compactness a finite subcovering  $\{U_{x_1}, \dots, U_{x_r}\}$  of  $\partial B$ . Let  $\ell_i := u_i^* L u_i \in \mathbb{R}[X]_1$  and

$$S := \{x \in \mathbb{R}^n \mid \ell_1(x) \geq 0, \dots, \ell_r(x) \geq 0\}.$$

Clearly,  $S_L \subseteq S$  and  $S \cap \partial B = \emptyset$ . Since  $S_L$  is non-empty by hypothesis and contained in  $B$ , it follows that  $S$  contains a point of  $B$ . But then it follows from the convexity of  $S$  together with  $S \cap \partial B = \emptyset$  that  $S \subseteq B$ . In particular,  $S$  is bounded.

Now every  $\ell_i \in C_L \subseteq M_L$ . Hence the quadratic module  $M$  generated by the  $\ell_i$  is contained in  $M_L$ . Choose  $N \in \mathbb{N}$  with  $N \pm X_i > 0$  on  $S$  for all  $i$ . Fix an  $i$  and  $\delta \in \{-1, 1\}$ . The system of linear inequalities

$$-N + \delta x_i \geq 0, \ell_1(x) \geq 0, \dots, \ell_r(x) \geq 0$$

is infeasible. Hence by Farkas' lemma [Fa], there are  $\alpha_j \in \mathbb{R}_{\geq 0}$  satisfying

$$-1 = \alpha_0(-N + \delta X_i) + \alpha_1 \ell_1 + \dots + \alpha_r \ell_r. \quad (3)$$

Note  $\alpha_0 \neq 0$  since  $S \neq \emptyset$ . Rearranging terms in (3) yields  $N - \delta X_i \in C_L$ . Since  $i$  and  $\delta$  were arbitrary and  $C_L \subseteq M_L$ , we conclude that  $M_L$  is archimedean.  $\blacksquare$

Surprisingly, establishing Lemma 2.1.1 for *empty* spectrahedra is more involved and will occupy us in the next subsection.

**2.2. Quadratic modules describing the empty spectrahedron.** The following is an extension of Farkas' lemma from LP to SDP due to Sturm [St, Lemma 2.18]. We include its simple proof based on a Hahn-Banach separation argument.

**Lemma 2.2.1** (Sturm). *A linear pencil  $L$  is strongly infeasible if and only if  $-1 \in C_L$ .*

*Proof.* Suppose

$$L = A_0 + \sum_{i=1}^n X_i A_i$$

is strongly infeasible,  $A_i \in S\mathbb{R}^{m \times m}$ . Then the non-empty convex sets  $\{L(x) \mid x \in \mathbb{R}^n\}$  and  $S\mathbb{R}_{\geq 0}^{m \times m}$  can be *strictly* separated by an affine hyperplane (since their Minkowski sums with a small ball are still disjoint and can therefore be separated [Ba2, Chapter III, Theorem 1.2]). This means that there is a non-zero linear form

$$\ell: S\mathbb{R}^{m \times m} \rightarrow \mathbb{R}$$

and  $\alpha \in \mathbb{R}$  with  $\ell(S\mathbb{R}_{\geq 0}^{m \times m}) \subseteq \mathbb{R}_{>\alpha}$  and  $\ell(\{L(x) \mid x \in \mathbb{R}^n\}) \subseteq \mathbb{R}_{<\alpha}$ . Choose  $B \in S\mathbb{R}^{m \times m}$  such that

$$\ell(A) = \text{tr}(AB)$$

for all  $A \in S\mathbb{R}^{m \times m}$ . Since  $\ell(S\mathbb{R}_{\geq 0}^{m \times m})$  is bounded from below, by the self-duality of the convex cone of positive semidefinite matrices,  $0 \neq B \succeq 0$ . Similarly, we obtain  $\ell(A_i) = 0$  for  $i \in \{1, \dots, n\}$ . Note that  $\alpha < 0$  since  $0 = \ell(0) \in \mathbb{R}_{>\alpha}$  so we can assume  $\ell(A_0) = -1$  by scaling. Writing  $B = \sum_i u_i u_i^*$  with  $u_i \in \mathbb{R}^m$ , we obtain

$$-1 = \ell(A_0) = \ell(L(x)) = \text{tr}(L(x) \sum_i u_i u_i^*) = \sum_i u_i^* L(x) u_i.$$

for all  $x \in \mathbb{R}^n$ . Hence  $-1 = \sum_i u_i^* L u_i \in C_L$ .

Conversely, if  $-1 \in C_L$ , i.e.,  $-1 = c + \sum_i u_i^* L u_i$  for some  $c \geq 0$  and  $u_i \in \mathbb{R}^m$ , then with  $B := \sum_i u_i u_i^* \in S\mathbb{R}_{\geq 0}^{m \times m}$  we obtain a linear form

$$\ell: S\mathbb{R}^{m \times m} \rightarrow \mathbb{R}, \quad A \mapsto \text{tr}(AB)$$

satisfying  $\ell(S\mathbb{R}_{\geq 0}^{m \times m}) \subseteq \mathbb{R}_{\geq 0}$  and  $\ell(\{L(x) \mid x \in \mathbb{R}^n\}) = \{-1 - c\} \subseteq \mathbb{R}_{\leq -1}$ . So  $L$  is strongly infeasible.  $\blacksquare$

**Lemma 2.2.2.** *Let  $L$  be an infeasible linear pencil of size  $m$ . The following are equivalent:*

- (i)  $L$  is weakly infeasible;
- (ii)  $S_{L+\varepsilon I_m} \neq \emptyset$  for all  $\varepsilon > 0$ .

*Proof.* Since all norms on a finite-dimensional vector space are equivalent, we can use the operator norm on  $\mathbb{R}^{m \times m}$ .

Suppose that (i) holds and  $\varepsilon > 0$  is given. Choose  $A \in S\mathbb{R}_{\geq 0}^{m \times m}$  and  $x \in \mathbb{R}^n$  with  $\|A - L(x)\| < \varepsilon$ . Then  $L(x) + \varepsilon I_m \succeq 0$ , i.e.,  $x \in S_{L+\varepsilon I_m}$ .

Conversely, suppose that (ii) holds. To show that

$$\text{dist}(\{L(x) \mid x \in \mathbb{R}^n\}, S\mathbb{R}_{\geq 0}^{m \times m}) = 0,$$

we let  $\varepsilon > 0$  be given and have to find  $A \in S\mathbb{R}_{\geq 0}^{m \times m}$  and  $x \in \mathbb{R}^n$  with

$$\|L(x) - A\| \leq \varepsilon.$$

But this is easy: choose  $x \in \mathbb{R}^n$  with  $L(x) + \varepsilon I_m \succeq 0$ , and set  $A := L(x) + \varepsilon I_m$ .  $\blacksquare$

The following lemma is due to Bohnenblust [Bo] (see also [Ba1, Theorem 4.2] for an easier accessible reference). While Bohnenblust gave a non-trivial bound on the number of terms that are really needed to test condition (i) below, we will not need this improvement and therefore take the trivial bound  $m$ . Then the proof becomes easy and we include it for the convenience of the reader.

**Lemma 2.2.3** (Bohnenblust). *For  $A_1, \dots, A_n \in S\mathbb{R}^{m \times m}$  the following are equivalent:*

- (i) Whenever  $u_1, \dots, u_m \in \mathbb{R}^m$  with  $\sum_{i=1}^m u_i^* A_j u_i = 0$  for all  $j \in \{1, \dots, n\}$ , then  $u_1 = \dots = u_m = 0$ ;
- (ii)  $\text{span}(A_1, \dots, A_n)$  contains a positive definite matrix.

*Proof.* It is trivial that (ii) implies (i). To prove that (i) implies (ii), note that  $S\mathbb{R}_{\geq 0}^{m \times m} = \{\sum_{i=1}^m u_i u_i^* \mid u_1, \dots, u_m \in \mathbb{R}^m\}$  and  $\sum_{i=1}^m u_i^* A u_i = \text{tr}(A \sum_{i=1}^m u_i u_i^*)$  for all  $u_1, \dots, u_m \in \mathbb{R}^m$ . The hypotheses thus say exactly that, given any  $B \in S\mathbb{R}_{\geq 0}^{m \times m}$ , we have

$$\text{tr}(A_1 B) = \dots = \text{tr}(A_n B) = 0 \implies B = 0. \quad (4)$$

Now suppose that  $\text{span}(A_1, \dots, A_n) \cap S\mathbb{R}_{\geq 0}^{m \times m} = \emptyset$ . By the standard separation theorem for two non-empty disjoint convex sets (see for example [Ba2, Chapter III, Theorem 1.2]),  $\text{span}(A_1, \dots, A_n)$  and  $S\mathbb{R}_{\geq 0}^{m \times m}$  can be separated by a hyperplane

(the separating affine hyperplane must obviously contain the origin). Therefore there is a non-zero linear form  $L: S\mathbb{R}^{m \times m} \rightarrow \mathbb{R}$  with

$$L(S\mathbb{R}_{>0}^{m \times m}) \subseteq \mathbb{R}_{\geq 0} \quad \text{and} \quad L(\text{span}(A_1, \dots, A_n)) \subseteq \mathbb{R}_{\leq 0}.$$

Then of course  $L(S\mathbb{R}_{\geq 0}^{m \times m}) \subseteq \mathbb{R}_{\geq 0}$  and  $L(\text{span}(A_1, \dots, A_n)) = \{0\}$ . Now choose  $B \in S\mathbb{R}^{m \times m}$  such that

$$L(A) = \text{tr}(AB) \quad \text{for all} \quad A \in S\mathbb{R}^{m \times m}.$$

Then  $B \neq 0$ ,  $B \in S\mathbb{R}_{\geq 0}^{m \times m}$  and  $\text{tr}(A_1 B) = \dots = \text{tr}(A_n B) = 0$ , contradicting (4). ■

**Lemma 2.2.4.** *Let  $L$  be a linear pencil of size  $m$  which is either weakly infeasible or weakly feasible. Then there are  $k \geq 1$  and  $u_1, \dots, u_k \in \mathbb{R}^m \setminus \{0\}$  such that  $\sum_{i=1}^k u_i^* L u_i = 0$ .*

*Proof.* Assume that the conclusion is not true. By Lemma 2.2.3 there are hence  $x_0, x_1, \dots, x_n \in \mathbb{R}$  such that

$$x_0 A_0 + x_1 A_1 + \dots + x_n A_n \succ 0.$$

Of course it is impossible that  $x_0 > 0$  since otherwise  $L(0) \succ 0$ . Also  $x_0 = 0$  is excluded (since otherwise  $L(cx_1, \dots, cx_n) \succ 0$  for  $c > 0$  large enough). Hence without loss of generality  $x_0 = -1$ , i.e.,  $x_1 A_1 + \dots + x_n A_n \succ A_0$ . Choose  $\varepsilon > 0$  such that

$$x_1 A_1 + \dots + x_n A_n \succ A_0 + 2\varepsilon I_m.$$

By Lemma 2.2.2, we can choose some  $y \in S_{L+\varepsilon I_m}$ . But then

$$A_0 + (x_1 + 2y_1)A_1 + \dots + (x_n + 2y_n)A_n \succ 2(A_0 + \varepsilon I_m + y_1 A_1 + \dots + y_n A_n) \succeq 0,$$

contradicting the hypotheses. ■

We come now to our first main theorem which is a version of Farkas' lemma for SDP which unlike Lemma 2.2.1 does not only work for strongly but also for weakly infeasible linear pencils. The price we pay is that we have to replace the finite-dimensional convex cone  $C_L$  by the infinite-dimensional quadratic module  $M_L$ . We will spend most of the sequel to refine and tame this result, cf. Theorems 2.3.1 and 3.5.2 below.

**Theorem 2.2.5.** *Let  $L$  be a linear pencil. Then*

$$S_L = \emptyset \quad \iff \quad -1 \in M_L.$$

*Proof.* One direction is trivial since  $M_L$  contains only polynomials nonnegative on  $S_L$ . We show the nontrivial implication from left to right by induction on the number  $n$  of variables appearing in  $L$ . In the case  $n = 0$ , we have  $L \in S\mathbb{R}^{m \times m}$  and therefore  $S_L = \emptyset \iff L \not\succeq 0 \iff -1 \in M_L$ .

For the induction step suppose  $n \geq 1$  and the statement is already known for linear pencils in less than  $n$  variables. The way we will later use the induction hypothesis is explained by the following claims which are weakening of the theorem itself.

Note that Claim 2 is just a special case of Claim 3' which we formulate for the sake of clarity. The proof of Claim 3' is very conceptual and uses a tool of real algebraic geometry, namely Prestel's semiorderings [PD]. For readers that are not familiar with this theory we prove a special case of it by an ad hoc technical argument. This special case of Claim 3' is stated in Claim 3 and suffices for our subsequent application.

**Claim 1.** If  $S_L = \emptyset$  and  $0 \neq \ell \in \mathbb{R}[X]_1$ , then  $-1 \in M_L + (\ell)$ .

*Explanation.* After an affine linear variable transformation, we may assume  $\ell = X_n$ . Set  $L' := L(X_1, \dots, X_{n-1}, 0)$ . Then  $-1 \in M_L + (X_n)$  is equivalent to

$-1 \in M_{L'}$ . By induction hypothesis, the latter follows if  $S_{L'} = \emptyset$ . But emptiness of  $S_L$  clearly implies the emptiness of  $S_{L'}$ .  $\square$

**Claim 2.** If  $S_L = \emptyset$  and  $0 \neq \ell \in M_L \cap -M_L$ , then  $-1 \in M_L$ .

*Explanation.* This follows from Claim 1 using the fact that  $M_L \cap -M_L$  is an ideal in  $\mathbb{R}[\underline{X}]$  which follows from (1).  $\square$

**Claim 3.** If  $S_L = \emptyset$  and  $0 \neq \ell \in \mathbb{R}[\underline{X}]_1$  such that  $-\ell^2 \in M_L$ , then  $-1 \in M_L$ .

*Explanation.* By Claim 1, there is  $p \in M_L$  and  $q \in \mathbb{R}[\underline{X}]$  such that  $-1 = p + q\ell$ . Now

$$\begin{aligned} -1 &= -2 + 1 = (2p + 2q\ell) + ((1 + q\ell)^2 - 2q\ell + q^2(-\ell^2)) \\ &= 2p + q^2(-\ell^2) + (1 + q\ell)^2 \in M_L, \end{aligned}$$

as desired.  $\square$

**Claim 3'.** If  $S_L = \emptyset$ ,  $k \geq 1$  and  $\ell_1, \dots, \ell_k \in \mathbb{R}[\underline{X}]_1$  such that  $0 \neq \ell_1 \cdots \ell_k \in M_L \cap -M_L$ , then  $-1 \in M_L$ .

*Explanation.* Assume  $-1 \notin M_L$ . Then there is a semiordering  $S$  of  $\mathbb{R}[\underline{X}]$  such that  $M_L \subseteq S$ . The advantage of  $S$  over  $M_L$  is that  $S \cap -S$  is always a prime ideal. Hence  $\ell_i \in S \cap -S$  for some  $i \in \{1, \dots, k\}$ . But this shows that  $-1 \notin M_L + (\ell_i)$  (even though  $(\ell_i)$  might not be contained in  $M_L \cap -M_L$ ). Hence  $S_L \neq \emptyset$  by Claim 1.  $\square$

Now we are prepared to do the induction step. Suppose  $S_L = \emptyset$ . If  $L$  is strongly infeasible, then  $-1 \in C_L \subseteq M_L$  by Lemma 2.2.1. Therefore we assume from now on that  $L$  is weakly infeasible. By Lemma 2.2.4, one of the following two cases applies:

**Case 1.** There is  $u \in \mathbb{R}^m \setminus \{0\}$  with  $u^*Lu = 0$ .

Of course, we can assume that  $L$  has no zero column since otherwise we can remove it together with the corresponding row without altering  $S_L$  and  $M_L$ . Write  $L = (\ell_{ij})_{1 \leq i, j \leq m}$ . By changing coordinates on  $\mathbb{R}^m$ , we can assume that  $u$  is the first standard basis vector, i.e.,  $\ell_{11} = 0$ . Moreover, we may assume  $\ell_{12} \neq 0$ . Then

$$2pq\ell_{12} + q^2\ell_{22} \in M_L$$

for all  $p, q \in \mathbb{R}[\underline{X}]$ . In particular,  $\ell_{12}^2(\ell_{22} + 2p) \in M_L$  for all  $p \in \mathbb{R}[\underline{X}]$  and therefore  $-\ell_{12}^2 \in M_L$ . Now we conclude by Claim 3 or 3' that  $-1 \in M_L$ .

**Case 2.** Case 1 does not apply but there are  $k \geq 2$  and  $u_1, \dots, u_k \in \mathbb{R}^m \setminus \{0\}$  such that  $\sum_{i=1}^k u_i^*Lu_i = 0$ .

In this case,

$$\ell := u_1^*Lu_1 = -\sum_{i=2}^k u_i^*Lu_i \in C_L \cap -C_L \subseteq M_L \cap -M_L$$

and we are done by Claim 2, since  $\ell = u_1^*Lu_1 \neq 0$ .  $\blacksquare$

**Corollary 2.2.6.** *Let  $L$  be a linear pencil. Then*

$$S_L \text{ is bounded} \iff M_L \text{ is archimedean.}$$

*Proof.* If  $S_L \neq \emptyset$ , then this is Lemma 2.1.1. If  $S_L = \emptyset$ , then  $-1 \in M_L$  by Theorem 2.2.5, so  $M_L$  is archimedean.  $\blacksquare$

**Remark 2.2.7.** Note that the above corollary is a strong variant of Schmüdgen's characterization [Sm] of bounded basic closed semialgebraic sets as being exactly those whose describing finitely generated preorderings are archimedean. Preorderings have the tendency of being much larger than quadratic modules. In general, a finitely generated quadratic module might describe a bounded or even an empty set without being archimedean, see [PD, Example 6.3.1] and [Ma, Example 7.3.2]. Corollary 2.2.6 says that quadratic modules associated to linear pencils behave very well in this respect.



We conclude this section with a version of Putinar's Positivstellensatz [Pu] for bounded spectrahedra:

**Corollary 2.2.8.** *Let  $L$  be a linear pencil and assume that  $S_L$  is bounded. If  $f \in \mathbb{R}[\underline{X}]$  satisfies  $f|_{S_L} > 0$ , then  $f \in M_L$ .*

*Proof.* By Corollary 2.2.6,  $M_L$  is archimedean. Now apply (a slight generalization of) Putinar's Positivstellensatz [Ma, Theorem 5.6.1].  $\blacksquare$

**Remark 2.2.9.** Let  $L$  be a linear pencil with bounded  $S_L$ .

- (1) In the case  $L$  is strongly feasible, Corollary 2.2.8 has already been proved in [HKM, §7] by completely different techniques, namely complete positivity from operator algebras. Note however that the more involved case in our approach occurs when  $L$  is infeasible.
- (2) From Corollary 8 it is easy to see that the quadratic module *in the sense of rings with involution* (see [KS]) associated to a linear pencil  $L$  of size  $m$  in the ring of  $s \times s$  matrix polynomials is archimedean (in the sense of [KS, Subsection 3.1] or [HKM, §6, §7]) if the spectrahedron  $S_L$  defined by  $L$  is bounded (cf. [HKM, §7]). Among other consequences, this implies a suitable generalization of Corollary 2.2.8 for *matrix polynomials* positive definite on the bounded spectrahedron  $S_L$  (cf. [HS, Corollary 1], [KS, Theorem 3.7] and [HKM, Theorem 7.1]).

**2.3. Degree Bounds.** Given a linear pencil  $L$  of size  $m$  and  $k \in \mathbb{N}_0$ , let

$$\begin{aligned} M_L^{(k)} &:= \left\{ \sum_i p_i^2 + \sum_j v_j^* L v_j \mid p_i \in \mathbb{R}[\underline{X}]_k, v_j \in \mathbb{R}[\underline{X}]_k^m \right\} \\ &= \left\{ s + \text{tr}(LS) \mid s \in \mathbb{R}[\underline{X}]_{2k} \text{ sos-polynomial, } S \in \mathbb{R}[\underline{X}]_{2k}^{m \times m} \text{ sos-matrix} \right\} \\ &\subseteq \mathbb{R}[\underline{X}]_{2k+1}, \end{aligned}$$

be the *truncated quadratic module with degree restriction  $k$*  associated to  $L$ . Note that  $M_L^{(0)} = C_L$ .

By a closer inspection of the proof of Theorem 2.2.5 we can obtain exponential degree bounds for the infeasibility certificate.

**Theorem 2.3.1.** *Let  $L$  be an infeasible linear pencil in  $n$  variables. Then*

$$-1 \in M_L^{(2^n - 1)}.$$

*Proof.* We shall prove this by induction on  $n$ . The statement is clear for  $n = 0$ . Given  $n \in \mathbb{N}$  and

$$L = A_0 + \sum_{i=1}^n X_i A_i,$$

we assume the statement has been established for all infeasible linear pencils with  $n - 1$  variables.

By Lemma 2.2.4, there is an  $u \in \mathbb{R}^{m^2} \setminus \{0\}$  with  $u^*(I_m \otimes L)u = 0$ . Replacing  $L$  by  $I_m \otimes L$  changes neither  $M_L^{(k)}$  nor  $S_L = \emptyset$ . Without loss of generality, we assume therefore that there is  $u \in \mathbb{R}^m \setminus \{0\}$  with  $u^* L u = 0$ . Writing  $L = (\ell_{ij})_{1 \leq i, j \leq m}$  and performing a linear coordinate change on  $\mathbb{R}^m$ , we can moreover assume  $\ell_{11} = 0$ . Furthermore, without loss of generality,  $\ell_{12} \neq 0$ . As in Case 1 of the proof of Theorem 2.2.5 above, we deduce

$$-\ell_{12}^2 \in M_L^{(1)}. \tag{5}$$

If  $\ell_{12} \in \mathbb{R}$ , we are done. Otherwise after possibly performing an affine linear change of variables on  $\mathbb{R}^n$ , we may assume  $\ell_{12} = X_n$ .

Now  $L' := L(X_1, \dots, X_{n-1}, 0)$  is an infeasible linear pencil in  $n-1$  variables. By our induction hypothesis,  $-1 \in M_{L'}^{(2^{n-1}-1)}$ . In particular, there are

$$p_i \in \mathbb{R}[\underline{X}]_{2^{n-1}-1} \text{ and } v_j \in \mathbb{R}[\underline{X}]_{2^{n-1}-1}^m$$

satisfying

$$-1 = \sum_i p_i^2 + \sum_j v_j^* L' v_j.$$

Let  $q := \sum_j v_j^* A_n v_j \in \mathbb{R}[\underline{X}]_{2^n-2}$ . Then

$$\begin{aligned} -1 &= 2 \sum_i p_i^2 + 2 \sum_j v_j^* L' v_j + 1 \\ &= 2 \sum_i p_i^2 + 2 \sum_j v_j^* L v_j - 2qX_n + 1 \\ &= 2 \sum_i p_i^2 + 2 \sum_j v_j^* L v_j + (1 - qX_n)^2 + q^2(-X_n^2). \end{aligned} \tag{6}$$

Since  $\deg q \leq 2^n - 2$ , we have  $q^2(-X_n) \in M_L^{(2^n-1)}$  by (5). Taken together with  $(1 - qX_n)^2 \in M_L^{(2^n-1)}$ , (6) implies  $-1 \in M_L^{(2^n-1)}$ .  $\blacksquare$

Similarly as in Theorem 2.3.1, one can obtain a bound on  $k$  in a certificate of the form  $-1 \in M_L^{(k)}$  that depends exponentially on the size  $m$  of  $L$ .

**Theorem 2.3.2.** *Let  $L$  be an infeasible linear pencil of size  $m$ . Then*

$$-1 \in M_L^{(2^{m-1}-1)}.$$

*Proof.* We prove this by induction on  $m$ . The statement is clear for  $m = 1$ . Given

$$L = A_0 + \sum_{i=1}^n X_i A_i \in S\mathbb{R}[\underline{X}]^{m \times m}$$

of size  $m \geq 2$ , we assume the statement has been established for all infeasible linear pencils of size  $m-1$ . If  $L$  is strongly infeasible, then  $-1 \in C_L = M_L^{(0)}$  by Lemma 2.2.1. So we may assume  $L$  is weakly infeasible.

**Claim.** There is an affine linear change of variables after which  $L$  assumes the form

$$L = \begin{bmatrix} b_0 & b^* \\ b & L' \end{bmatrix},$$

where  $b_0 \in \mathbb{R}[\underline{X}]_1$ ,  $b = [b_1 \ \dots \ b_{m-1}]^* \in \mathbb{R}[\underline{X}]_1^{m-1}$ ,  $L'$  is a linear pencil of size  $m-1$ , and  $b_j \in \mathbb{R}[\underline{X}]_1$  satisfy

$$-b_j^2 \in M_L^{(1)} \quad \text{for } j = 0, \dots, m-1. \tag{7}$$

Furthermore,  $b_0$  can be chosen to be either 0 or  $X_1$ .

*Explanation.* By Lemma 2.2.4, there is  $k \in \mathbb{N}$  and  $u_1, \dots, u_k \in \mathbb{R}^m \setminus \{0\}$  with  $\sum_{i=1}^k u_i^* L u_i = 0$ . We distinguish two cases.

**Case 1.** There is  $u \in \mathbb{R}^m \setminus \{0\}$  with  $u^* L u = 0$ .

Write  $L = (\ell_{ij})_{1 \leq i, j \leq m}$ . By changing coordinates on  $\mathbb{R}^m$ , we can assume that  $u$  is the first standard basis vector, i.e.,  $\ell_{11} = 0$ . Hence

$$L = \begin{bmatrix} 0 & b^* \\ b & L' \end{bmatrix},$$

where  $b = [b_1 \ \dots \ b_{m-1}]^* \in \mathbb{R}[\underline{X}]_1^{m-1}$  and  $L'$  is a linear pencil of size  $m-1$ . As in the proof of Theorem 2.3.1, we deduce  $-b_j^2 \in M_L^{(1)}$  for all  $j = 1, \dots, m-1$ .

**Case 2.** Case 1 does not apply but there are  $k \geq 2$  and  $u_1, \dots, u_k \in \mathbb{R}^m \setminus \{0\}$  such that  $\sum_{i=1}^k u_i^* L u_i = 0$ .

In this case,

$$\ell_{11} := u_1^* L u_1 = - \sum_{i=2}^k u_i^* L u_i \in C_L \cap -C_L = M_L^{(0)} \cap -M_L^{(0)}.$$

Since Case 1 does not apply,  $\ell_{11} \neq 0$ . Furthermore, since  $L$  is assumed to be weakly infeasible,  $\ell_{11} \notin \mathbb{R}$ . Hence after an affine linear change of variables on  $\mathbb{R}^n$ , we can assume  $\ell_{11} = X_1$ . Thus

$$L = \begin{bmatrix} X_1 & b^* \\ b & L' \end{bmatrix},$$

where  $b = [b_1 \ \dots \ b_{m-1}]^* \in \mathbb{R}[\underline{X}]_1^{m-1}$  and  $L'$  is a linear pencil of size  $m-1$ . Note that

$$-4X_1^2 = (1 - X_1)^2 X_1 + (1 + X_1)^2 (-X_1)$$

shows that  $-X_1^2 \in M_L^{(1)}$ . Using this, one gets similarly as above that also each of the entries  $b_j$  of  $b$  satisfies  $-b_j^2 \in M_L^{(1)}$ . This proves our claim.  $\square$

If one of the  $b_j \in \mathbb{R} \setminus \{0\}$ , we are done by (7). Otherwise we consider two cases.

**Case a.** If the linear system  $b_0(x) = 0, b(x) = 0$  is infeasible, then we proceed as follows. There are  $\alpha_0, \dots, \alpha_{m-1} \in \mathbb{R}$  satisfying

$$\sum_{j=0}^{m-1} \alpha_j b_j = 1. \quad (8)$$

For each  $j = 0, \dots, m-1$  and  $\delta \in \mathbb{R}$  we have

$$1 + \delta b_j = \left(1 + \frac{\delta}{2} b_j\right)^2 + \frac{\delta^2}{4} (-b_j^2) \in M_L^{(1)}$$

by (7). Hence (8) implies

$$-1 = 1 - 2 = 1 - 2 \sum_{j=0}^{m-1} \alpha_j b_j = \sum_{j=0}^{m-1} \left(\frac{1}{m} - 2\alpha_j b_j\right) \in M_L^{(1)}.$$

**Case b.** Suppose the linear system  $b_0(x) = b(x) = 0$  is feasible. Then we perform an affine linear change of variables on  $\mathbb{R}^n$  to ensure

$$\{x \in \mathbb{R}^n \mid b_0(x) = 0, b(x) = 0\} = \{0\}^r \times \mathbb{R}^{n-r}$$

for some  $r \in \mathbb{N}$ . Moreover, we may assume  $X_1, \dots, X_r$  are among the entries  $b_j$ ,  $j = 0, \dots, m-1$ .

Now  $L'' := L'(0, \dots, 0, X_{r+1}, \dots, X_n)$  is an infeasible linear pencil of size  $m-1$ . By our induction hypothesis,  $-1 \in M_{L''}^{(2^{m-2}-1)}$ . In particular, there are  $s \in \sum \mathbb{R}[\underline{X}]^2$  with  $\deg s \leq 2^{m-1} - 2$ , and  $v_i \in \mathbb{R}[\underline{X}]_{2^{m-2}-1}^{m-1}$  satisfying

$$-1 = s + \sum_i v_i^* L'' v_i.$$

Introducing

$$q_t := \sum_i v_i^* A_t v_i \in \mathbb{R}[\underline{X}]_{2^{m-1}-2} \quad \text{and} \quad w_i := \begin{bmatrix} 0 \\ v_i \end{bmatrix} \in \mathbb{R}[\underline{X}]_{2^{m-2}-1}^m$$

we have

$$\begin{aligned}
-1 &= (2s + 2 \sum_i v_i^* L'' v_i) + 1 \\
&= (2s + 2 \sum_i v_i^* L' v_i - \sum_{t=1}^r 2q_t X_t) + \sum_{t=1}^r \left( \left( \frac{1}{\sqrt{r}} - \sqrt{r} q_t X_t \right)^2 + 2q_t X_t + r q_t^2 (-X_t^2) \right) \\
&= 2s + 2 \sum_i w_i^* L w_i + \sum_{t=1}^r \left( \frac{1}{\sqrt{r}} - \sqrt{r} q_t X_t \right)^2 + \sum_{t=1}^r r q_t^2 (-X_t^2).
\end{aligned} \tag{9}$$

Combining  $q_t^2(-X_t^2) \in M_L^{(2^{m-1}-1)}$  with  $(\frac{1}{\sqrt{r}} - \sqrt{r} q_t X_t)^2 \in M_L^{(2^{m-1}-1)}$ , (9) implies  $-1 \in M_L^{(2^{m-1}-1)}$ .  $\blacksquare$

**Corollary 2.3.3.** *Let  $L$  be an infeasible linear pencil of size  $m$  in  $n$  variables. Then*

$$-1 \in M_L^{(2^{\min\{m-1, n\}}-1)}.$$

*Proof.* This is immediate from Theorems 2.3.1 and 2.3.2.  $\blacksquare$

**2.4. Examples.** The standard textbook example [St, WSV] of a weakly infeasible linear pencil seems to be

$$L := \begin{bmatrix} X & 1 \\ 1 & 0 \end{bmatrix}.$$

Then  $-1 \notin C_L$ , but  $-1 \in M_L^{(1)}$ . Indeed, for

$$u := \left[ 1 \quad -1 - \frac{X}{2} \right]^*,$$

we have

$$-1 = \frac{1}{2} u^* L u.$$

**Example 2.4.1.** Let

$$L := \begin{bmatrix} 0 & X_1 & 0 \\ X_1 & X_2 & 1 \\ 0 & 1 & X_1 \end{bmatrix}.$$

Then  $L$  is weakly infeasible and  $-1 \notin M_L^{(1)}$ .

Assume otherwise, and let

$$-1 = s + \sum_j v_j^* L v_j, \tag{10}$$

where  $v_j \in \mathbb{R}[\underline{X}]_1^3$  and  $s \in \sum \mathbb{R}[\underline{X}]^2$  with  $\deg s \leq 2$ . We shall carefully analyze the terms  $v_j^* L v_j$ . Write

$$v_j = [q_{1j} \quad q_{2j} \quad q_{3j}]^*, \quad \text{and} \quad q_{ij} = a_{ij} + b_{ij} X_1 + c_{ij} X_2$$

with  $a_{ij}, b_{ij}, c_{ij} \in \mathbb{R}$ . Then the  $X_2^3$  coefficient of  $v_j^* L v_j$  equals  $c_{2j}^2$ , so  $c_{2j} = 0$  for all  $j$ . Next, by considering the  $X_1 X_2^2$  terms, we deduce  $c_{3j} = 0$ . Now the only terms possibly contributing to  $X_2^2$  come from  $s$ , so  $s \in \mathbb{R}[\underline{X}]_2$ . The coefficient of  $X_2$  in  $v_j^* L v_j$  is a square, so  $a_{2j} = 0$ . But now  $v_j^* L v_j$  does not have a constant term anymore, leading to a contradiction with (10).

From Theorem 2.2.5 it follows that  $-1 \in M_L^{(3)}$ . In fact,  $-1 \in M_L^{(2)}$  since

$$-2 = u^* L u$$

for  $u = \left[ \frac{1}{2} + \frac{X_2}{2} + \frac{X_2^2}{8} \quad -1 \quad 1 + \frac{X_2}{2} \right]^* \in \mathbb{R}[\underline{X}]_2^3$ .

## 3. POLYNOMIAL SIZE CERTIFICATES FOR INFEASIBILITY

Theorem 2.3.1 (or Corollary 2.3.3) enables us to reformulate the feasibility of a linear pencil as the infeasibility of an LMI. Indeed, let a linear pencil  $L$  in  $n$  variables of size  $m$  be given. Then  $L$  is infeasible if and only if there exists an sos-polynomial  $s \in \mathbb{R}[\underline{X}]$  and an sos-matrix  $S \in \mathbb{R}[\underline{X}]^{m \times m}$  both of degree at most  $2^{n+1} - 2$  such that

$$-1 = s + \text{tr}(LS). \quad (11)$$

By comparing coefficients, the polynomial equation (11) can of course be written as a system of linear equations in the coefficients of  $s$  and  $S$ . Now (the coefficient tuples of) sos-polynomials in  $\mathbb{R}[\underline{X}]$  of bounded degree form a *projection* of a spectrahedron. In other words, the condition of being (the coefficient tuple) of an sos-polynomial in  $\mathbb{R}[\underline{X}]$  of bounded degree can be expressed within an LMI by means of additional variables. This is the well known *Gram matrix method* [Lau, Ma]. As noted by Kojima [Ko] and nicely described by Hol and Scherer [HS], the Gram matrix method extends easily to sos-matrices.

Hence the existence of  $s$  and  $S$  in (11) is indeed equivalent to the feasibility of an LMI which we will not write down explicitly. The drawback of this LMI is that it will be large since our degree bounds for the sos-polynomial and the sos-matrix are exponential.

This problem is overcome in this section, where for a given linear pencil  $L$ , we construct an LMI whose feasibility is equivalent to the infeasibility of the given linear pencil  $L$  and which can be written down in polynomial time (and hence has polynomial size) in the bit size of  $L$  if  $L$  has rational coefficients. Each feasible point of the new LMI gives rise to a certificate of infeasibility for  $L$  which is however more involved than the one using the quadratic module  $M_L$ . In addition to the quadratic module, we will now use another notion from real algebraic geometry, namely the real radical ideal.

But actually we will not only characterize infeasibility of LMIs but even give a new duality theory for SDP where strong duality always holds in contrast to the standard duality theory. We will call our dual the *sums of squares dual* of an SDP. For a given (primal) SDP with rational coefficients the sums of squares dual can be written down in polynomial time in the bit size of the primal.

**3.1. Review of standard SDP duality.** We first recall briefly the standard duality theory of SDP. We present it from the view point of a real algebraic geometer, i.e., we use the language of polynomials in the formulation of the primal-dual pair of SDPs and in the proof of strong duality. This is necessary for a good understanding of the sums of squares dual which we will give later.

A semidefinite program (P) and its standard dual (D) is given by a linear pencil  $L \in \mathbb{R}[\underline{X}]^{m \times m}$  and a linear polynomial  $\ell \in \mathbb{R}[\underline{X}]$  as follows:

$$\begin{array}{ll} (P) & \text{minimize } \ell(x) \\ & \text{subject to } x \in \mathbb{R}^n \\ & L(x) \succeq 0 \end{array} \quad \begin{array}{ll} (D) & \text{maximize } a \\ & \text{subject to } S \in S\mathbb{R}^{m \times m}, a \in \mathbb{R} \\ & S \succeq 0 \\ & \ell - a = \text{tr}(LS) \end{array}$$

To see that this corresponds (up to some minor technicalities) to the formulation in the literature, just write the polynomial constraint  $\ell - a = \text{tr}(LS)$  of the dual as  $n + 1$  linear equations by comparing coefficients.

The optimal values of (P) and (D) are defined to be

$$\begin{aligned} P^* &:= \inf\{\ell(x) \mid x \in \mathbb{R}^n, L(x) \succeq 0\} \in \mathbb{R} \cup \{\pm\infty\} \quad \text{and} \\ D^* &:= \sup\{a \mid S \in S\mathbb{R}_{\succeq 0}^{m \times m}, a \in \mathbb{R}, \ell - a = \text{tr}(LS)\} \in \mathbb{R} \cup \{\pm\infty\}, \end{aligned}$$

respectively, where the infimum and the supremum is taken in the ordered set  $\{-\infty\} \cup \mathbb{R} \cup \{\infty\}$  (where  $\inf \emptyset = \infty$  and  $\sup \emptyset = -\infty$ ).

By *weak duality*, we mean that  $P^* \geq D^*$  or equivalently that the objective value of  $(P)$  at any of its feasible points is greater or equal to the objective value of  $(D)$  at any of its feasible points.

Fix a linear pencil  $L$ . It is easy to see that weak duality holds for all primal objectives  $\ell$  if and only if

$$f \in C_L \implies f \geq 0 \text{ on } S_L$$

holds for all  $f \in \mathbb{R}[\underline{X}]_1$ , which is of course true. By *strong duality*, we mean that  $P^* = D^*$  (*zero duality gap*) and that (the objective of)  $(D)$  *attains* this common optimal value in case this is finite. It is a little exercise to see that strong duality for all primal objectives  $\ell$  is equivalent to

$$f \geq 0 \text{ on } S_L \iff f \in C_L$$

for all  $f \in \mathbb{R}[\underline{X}]_1$ .

As we have seen in Subsection 2.4, this fails in general. It is however well-known that it is true when the feasible set  $S_L$  of the primal  $(P)$  has non-empty interior (e.g. if  $L$  is strongly feasible). For convenience of the reader, we include a proof which has a bit of a flavor of real algebraic geometry.

**Proposition 3.1.1** (Standard SDP duality). *Let  $L \in S\mathbb{R}[\underline{X}]^{m \times m}$  be a linear pencil such that  $S_L$  has non-empty interior. Then*

$$f \geq 0 \text{ on } S_L \iff f \in C_L$$

for all  $f \in \mathbb{R}[\underline{X}]_1$ .

*Proof.* In a preliminary step, we show that the convex cone  $C_L$  is closed in  $\mathbb{R}[\underline{X}]_1$ . To this end, consider the linear subspace  $U := \{u \in \mathbb{R}^m \mid Lu = 0\} \subseteq \mathbb{R}^m$ . The map

$$\varphi: \mathbb{R} \times (\mathbb{R}^m/U)^m \rightarrow C_L, \quad (a, \bar{u}_1, \dots, \bar{u}_m) \rightarrow a^2 + \sum_{i=1}^m u_i^* L u_i$$

is well-defined and surjective.

Suppose  $\varphi$  maps  $(a, \bar{u}_1, \dots, \bar{u}_m) \in (\mathbb{R}^m/U)^m$  to 0. Fix  $i \in \{1, \dots, m\}$ . Then  $u_i^* L(x) u_i = 0$  for all  $x \in S_L$ . Since  $L(x) \succeq 0$ , this implies  $L(x) u_i = 0$  for all  $x \in S_L$ . Using the hypothesis that  $S_L$  has non-empty interior, we conclude that  $Lu_i = 0$ , i.e.,  $u_i \in U$ . Since  $i$  was arbitrary and  $a = 0$ , this yields  $(a, \bar{u}_1, \dots, \bar{u}_m) = 0$ .

This shows  $\varphi^{-1}(0) = \{0\}$ . Together with the fact that  $\varphi$  is a (quadratically) homogeneous map, this implies that  $\varphi$  is proper (see for example [PS, Lemma 2.7]). In particular,  $C_L = \text{im } \varphi$  is closed.

Suppose now that  $f \notin \mathbb{R}[\underline{X}]_1 \setminus C_L$ . The task is to find  $x \in S_L$  such that  $f(x) < 0$ . Being a closed convex cone,  $C_L$  is the intersection of all closed half-spaces containing it. Therefore we find a linear map  $\psi: \mathbb{R}[\underline{X}]_1 \rightarrow \mathbb{R}$  such that  $\psi(C_L) \subseteq \mathbb{R}_{\geq 0}$  and  $\psi(f) < 0$ . We can assume  $\psi(1) > 0$  since otherwise  $\psi(1) = 0$  and we can replace  $\psi$  by  $\psi + \varepsilon \text{ev}_y$  for some small  $\varepsilon > 0$  where  $y \in S_L$  is chosen arbitrarily. Hereby  $\text{ev}_x: \mathbb{R}[\underline{X}]_1 \rightarrow \mathbb{R}$  denotes the evaluation in  $x \in \mathbb{R}^n$ . Finally, after a suitable scaling we can even assume  $\psi(1) = 1$ .

Now setting  $x := (\psi(X_1), \dots, \psi(X_n)) \in \mathbb{R}^n$ , we have  $\psi = \text{ev}_x$ . So  $\psi(C_L) \subseteq \mathbb{R}_{\geq 0}$  means exactly that  $L(x) \succeq 0$ , i.e.,  $x \in S_L$ . At the same time  $f(x) = \psi(f) < 0$  as desired.  $\blacksquare$

**3.2. Certificates for low-dimensionality of spectrahedra.** As mentioned above, the problems with the standard duality theory for SDP arise when one deals with spectrahedra having empty interior. Every convex set with empty interior is contained in an affine hyperplane. The basic idea is now to code the search for such

an affine hyperplane into the dual SDP and to replace equality in the constraint  $f - a = \text{tr}(LS)$  of (D) by congruence modulo the linear polynomial  $\ell$  defining the affine hyperplane. This raises however several issues:

First,  $S_L$  might have codimension bigger than one in  $\mathbb{R}^n$ . This will be resolved by iterating the search up to  $n$  times.

Second, we do not see any possibility to encode the search for the linear polynomial  $\ell$  directly into an SDP. What we can implement is the search for a non-zero *quadratic* sos-polynomial  $q$  together with a certificate of  $S_L \subseteq \{q = 0\}$ . Note that  $\{q = 0\}$  is a proper affine subspace of  $\mathbb{R}^n$ . It would be ideal to find  $q$  such that  $\{q = 0\}$  is the affine hull of  $S_L$  since then we could actually avoid the  $n$ -fold iteration just mentioned. However it will follow from Example 3.2.2 below that this is in general not possible.

Third, we have to think about how to implement congruence modulo linear polynomials  $\ell$  vanishing on  $\{q = 0\}$ . This will be dealt with by using the radical ideal from real algebraic geometry in connection with Schur complements.

We begin with a result which ensures that a suitable quadratic sos-polynomial  $q$  can always be found. In fact, the following proposition says that there exists such a  $q$  which is actually a square. The statement is of interest in itself since it provides certificates for low-dimensionality of spectrahedra. We need *quadratic* (i.e., degree  $\leq 2$ ) sos-matrices for this.

**Proposition 3.2.1.** *For any linear pencil  $L \in S\mathbb{R}[\underline{X}]^{m \times m}$ , the following are equivalent:*

- (i)  $S_L$  has empty interior;
- (ii) There exists a non-zero linear polynomial  $\ell \in \mathbb{R}[\underline{X}]_1$  and a quadratic sos-matrix  $S \in S\mathbb{R}[\underline{X}]^{m \times m}$  such that

$$-\ell^2 = \text{tr}(LS). \quad (12)$$

*Proof.* From (ii) it follows that  $-\ell^2 \in M_L$  and therefore  $-\ell^2 \geq 0$  on  $S_L$ , which implies  $\ell = 0$  on  $S_L$ . So it is trivial that (ii) implies (i).

For the converse, suppose that  $S_L$  has empty interior. If there is  $u \in \mathbb{R}^m \setminus \{0\}$  such that  $Lu = 0$  then, by an orthogonal change of coordinates on  $\mathbb{R}^m$ , we could assume that  $u$  is the first unit vector  $e_1$ . But then we delete the first column and the first row from  $L$ . We can iterate this and therefore assume from now on that there is no  $u \in \mathbb{R}^m \setminus \{0\}$  with  $Lu = 0$ .

We first treat the rather trivial case where  $L$  is strongly infeasible. By Lemma 2.2.1, there are  $c \in \mathbb{R}_{\geq 0}$  and  $u_i \in \mathbb{R}^m$  with  $-1 - c = \sum_i u_i^* L u_i$ . By scaling the  $u_i$  we can assume  $c = 0$ . Setting  $S := \sum_i u_i u_i^* \in S\mathbb{R}^{m \times m}$  and  $\ell := 1$ , we have  $-\ell^2 = -1 = \sum_i u_i^* L u_i = \text{tr}(LS)$  for the constant sos-matrix  $S$  and the constant non-zero linear polynomial  $\ell$ .

Now we assume that  $L$  is weakly infeasible or feasible. In case that  $L$  is feasible, it is clearly weakly feasible since otherwise  $S_L$  would have non-empty interior. Now Lemma 2.2.4 justifies the following case distinction:

**Case 1.** There is  $u \in \mathbb{R}^m \setminus \{0\}$  with  $u^* L u = 0$ .

Write  $L = (\ell_{ij})_{1 \leq i, j \leq m}$ . Again by an orthogonal change of coordinates on  $\mathbb{R}^m$ , we can assume that  $u = e_1$ , i.e.,  $\ell_{11} = 0$ . Moreover, we may assume  $\ell := \ell_{12} \neq 0$  (since  $Le_1 = Lu \neq 0$ ). Setting  $\ell' := \frac{1}{2}(-1 - \ell_{22})$ ,  $v := [\ell' \ \ell \ 0 \dots 0]^*$  and  $S := vv^*$ , we have

$$\text{tr}(LS) = v^* L v = 2\ell' \ell \ell_{12} + \ell^2 \ell_{22} = \ell^2 (\ell_{22} + 2\ell') = -\ell^2.$$

**Case 2.** Case 1 does not apply but there are  $k \geq 2$  and  $u_1, \dots, u_k \in \mathbb{R}^m \setminus \{0\}$  such that  $\sum_{i=1}^k u_i^* L u_i = 0$ .

Here we set  $\ell := u_1^* L u_1 \neq 0$  and write  $-\ell = \ell_1^2 - \ell_2^2$  where  $\ell_1 := \frac{1}{2}(-\ell + 1) \in \mathbb{R}[\underline{X}]_1$  and  $\ell_2 := \frac{1}{2}(-\ell - 1) \in \mathbb{R}[\underline{X}]_1$ . Then we can use the quadratic sos-matrix

$$S := \ell_1^2 u_1 u_1^* + \ell_2^2 \sum_{i=2}^k u_i u_i^* = \ell_1^2 u_1 u_1^* - \ell_2^2 u_1 u_1^* = -\ell u_1 u_1^*$$

to get  $\text{tr}(LS) = -\ell u_1^* L u_1 = -\ell^2$ .  $\blacksquare$

The certificate (12) of low-dimensionality exists for *some* but in general *not for every* affine hyperplane containing the spectrahedron. We illustrate this by the following example where the spectrahedron has codimension two and therefore is contained in infinitely many affine hyperplanes *only one* of which allows for a certificate (12).

**Example 3.2.2.** Let

$$L = \begin{bmatrix} 0 & X_1 & 0 \\ X_1 & X_2 & X_3 \\ 0 & X_3 & X_1 \end{bmatrix}.$$

Then  $S_L = \{(0, x_2, 0) \in \mathbb{R}^3 \mid x_2 \geq 0\}$  and the (affine) hyperplanes containing  $S_L$  are  $\{X_1 = 0\}$  and  $\{aX_1 + X_3 = 0\}$  ( $a \in \mathbb{R}$ ). As is shown in Case 1 of the proof of Proposition 3.2.1, the certificate of low-dimensionality (12) exists for the hyperplane  $\{X_1 = 0\}$ , i.e., there is a quadratic sos-matrix  $S$  such that  $-X_1^2 = \text{tr}(LS)$ . However, none of the other hyperplanes containing  $S_L$  allows for the certificate (12).

Otherwise assume that there is  $a \in \mathbb{R}$  such that  $\{aX_1 + X_3 = 0\}$  has a also corresponding certificate. Combining it with the one for  $\{X_1 = 0\}$ , we get a quadratic sos-matrix  $S$  such that

$$-(2a^2)X_1^2 - 2(aX_1 + X_3)^2 = \text{tr}(LS)$$

which implies

$$-X_3^2 = (2aX_1 + X_3)^2 + (-(2a^2)X_1^2 - 2(aX_1 + X_3)^2) \in M_L^{(1)}.$$

Specializing  $X_3$  to 1, one gets the contradiction  $-1 \in M_{L'}^{(1)}$  where  $L'$  is the linear pencil from Example 2.4.1.

**3.3. Iterating the search for an affine hyperplane.** We now carry out the slightly technical but easy iteration of Proposition 3.2.1 announced in Subsection 3.2 and combine it with Proposition 3.1.1. We get a new type of Positivstellensatz for linear polynomials on spectrahedra with bounded degree complexity.

**Theorem 3.3.1** (Positivstellensatz for linear polynomials on spectrahedra).

Let  $L \in S\mathbb{R}[\underline{X}]^{m \times m}$  be a linear pencil and  $f \in \mathbb{R}[\underline{X}]_1$ . Then

$$f \geq 0 \text{ on } S_L$$

if and only if there exist  $\ell_1, \dots, \ell_n \in \mathbb{R}[\underline{X}]_1$ , quadratic sos-matrices  $S_1, \dots, S_n \in S\mathbb{R}[\underline{X}]^{m \times m}$ , a matrix  $S \in S\mathbb{R}_{\geq 0}^{m \times m}$  and  $c \geq 0$  such that

$$\ell_i^2 + \text{tr}(LS_i) \in (\ell_1, \dots, \ell_{i-1}) \quad \text{for } i \in \{1, \dots, n\}, \quad \text{and} \quad (13)$$

$$f - c - \text{tr}(LS) \in (\ell_1, \dots, \ell_n). \quad (14)$$

*Proof.* We first prove that  $f \geq 0$  on  $S_L$  in the presence of (13) and (14).

The traces in (13) and (14) are elements of  $M_L$  and therefore nonnegative on  $S_L$ . Hence it is clear that constraint (14) gives  $f \geq 0$  on  $S_L$  if we show that  $\ell_i$  vanishes on  $S_L$  for all  $i \in \{1, \dots, n\}$ . Fix  $i \in \{1, \dots, n\}$  and assume by induction that  $\ell_1, \dots, \ell_{i-1}$  vanish on  $S_L$ . Then (13) implies  $\ell_i^2 + \text{tr}(LS_i)$  vanishes on  $S_L$  and therefore also  $\ell_i$ .



Conversely, suppose now that  $f \geq 0$  on  $S_L$ . We will obtain the data with properties (13) and (14) by induction on the number of variables  $n \in \mathbb{N}_0$ .

To do the induction basis, suppose first that  $n = 0$ . Then  $f \geq 0$  on  $S_L$  just means that the real number  $f$  is nonnegative if  $L \in S\mathbb{R}^{m \times m}$  is positive semidefinite. But if  $f \geq 0$ , then it suffices to choose  $c := f \geq 0$  and  $S := 0$  to obtain (14) with  $n = 0$ , and the other condition (13) is empty since  $n = 0$ . We now assume that  $f < 0$  and therefore  $L \not\geq 0$ . Then we choose  $u \in \mathbb{R}^m$  with  $u^*Lu = f$ . Setting  $S := uu^* \in S\mathbb{R}_{\geq 0}^{m \times m}$  and  $c := 0$ , we have

$$f - c - \text{tr}(LS) = f - u^*Lu = f - f = 0,$$

as required.

For the induction step, we now suppose that  $n \in \mathbb{N}$  and that we know already how to find the required data for linear pencils in  $n - 1$  variables. We distinguish two cases and will use the induction hypothesis only in the second one.

**Case 1.**  $S_L$  contains an interior point.

In this case, we set all  $\ell_i$  and  $S_i$  to zero so that (13) is trivially satisfied. Property (14) can be fulfilled by Proposition 3.1.1.

**Case 2.** The interior of  $S_L$  is empty.

In this case, we apply Proposition 3.2.1 to obtain  $0 \neq \ell_1 \in \mathbb{R}[\underline{X}]_1$  and a quadratic sos-matrix  $S_1 \in S\mathbb{R}[\underline{X}]^{m \times m}$  with

$$\ell_1^2 + \text{tr}(LS_1) = 0. \quad (15)$$

The case where  $\ell_1$  is constant is trivial. In fact, in this case we can choose all remaining data being zero since  $(\ell_1, \dots, \ell_i) = (\ell_1) = \mathbb{R}[\underline{X}]$  for all  $i \in \{1, \dots, n\}$ .

From now on we therefore assume  $\ell_1$  to be non-constant. But then the reader easily checks that there is no harm carrying out an affine linear variable transformation which allows us to assume  $\ell_1 = X_n$ . We then apply the induction hypothesis to the linear pencil  $L' := L(X_1, \dots, X_{n-1}, 0)$  and the linear polynomial  $f' := f(X_1, \dots, X_{n-1}, 0)$  in  $n - 1$  variables to obtain  $\ell_2, \dots, \ell_n \in \mathbb{R}[\underline{X}]$ , quadratic sos-matrices  $S_2, \dots, S_n \in S\mathbb{R}[\underline{X}]^{m \times m}$ , a matrix  $S \in S\mathbb{R}_{\geq 0}^{m \times m}$  and a constant  $c \geq 0$  such that

$$\ell_i^2 + \text{tr}(L'S_i) \in (\ell_2, \dots, \ell_{i-1}) \quad \text{for } i \in \{2, \dots, n\} \quad \text{and} \quad (16)$$

$$f' - c - \text{tr}(L'S) \in (\ell_2, \dots, \ell_n). \quad (17)$$

Noting that both  $f - f'$  and  $\text{tr}(LS_i) - \text{tr}(L'S_i) = \text{tr}((L - L')S_i)$  are contained in the ideal  $(X_n) = (\ell_1)$ , we see that (16) together with (15) implies (13). In the same manner, (17) yields (14).  $\blacksquare$

**3.4. The real radical and Schur complements.** Let  $L \in S\mathbb{R}[\underline{X}]^{m \times m}$  be a linear pencil and  $q \in \mathbb{R}[\underline{X}]$  a (quadratic) sos-polynomial such that  $-q = \text{tr}(LS)$  for some (quadratic) sos-matrix  $S$  like in (12) above. In order to resolve the third issue mentioned in Subsection 3.2, we would like to get our hands on (cubic) polynomials vanishing on  $\{q = 0\}$ . In other words, we want to implement the ideals appearing in (13) and (14) in an SDP.

Recall that for any ideal  $\mathcal{I} \subseteq \mathbb{R}[\underline{X}]$ , its *radical*  $\sqrt{\mathcal{I}}$  and its *real radical*  $\sqrt[\mathbb{R}]{\mathcal{I}}$  are the ideals defined by

$$\begin{aligned} \sqrt{\mathcal{I}} &:= \{p \in \mathbb{R}[\underline{X}] \mid \exists k \in \mathbb{N} : f^k \in \mathcal{I}\} \quad \text{and} \\ \sqrt[\mathbb{R}]{\mathcal{I}} &:= \{p \in \mathbb{R}[\underline{X}] \mid \exists k \in \mathbb{N} : \exists s \in \sum \mathbb{R}[\underline{X}]^2 : f^{2k} + s \in \mathcal{I}\}. \end{aligned}$$

An ideal  $\mathcal{I} \subseteq \mathbb{R}[\underline{X}]$  is called *radical* if  $\mathcal{I} = \sqrt{\mathcal{I}}$  and *real radical* if  $\mathcal{I} = \sqrt[\mathbb{R}]{\mathcal{I}}$ .

By the Real Nullstellensatz [BCR, Ma, PD], each polynomial vanishing on the real zero set  $\{q = 0\}$  of  $q$  lies in  $\sqrt[\mathbb{R}]{(q)}$ . This gives a strategy of how to find the

cubic polynomials vanishing on  $\{q = 0\}$ , cf. Proposition 3.4.1 and Lemma 3.4.2 below. The Real Nullstellensatz plays only a motivating role for us. We will only use its trivial converse: Each element of  $\sqrt[q]{\mathbb{R}}$  vanishes on  $\{q = 0\}$ .

The question is now how to model the search for elements in the real radical ideal by SDP. The key to this will be to represent polynomials by matrices as it is done in the Gram matrix method mentioned at the beginning of Section 3. For this we have to introduce notation.

For each  $d \in \mathbb{N}_0$ , let  $s(d) := \dim \mathbb{R}[\underline{X}]_d = \binom{d+n}{n}$  denote the number of monomials of degree at most  $d$  in  $n$  variables and  $\overrightarrow{[\underline{X}]}_d \in \mathbb{R}[\underline{X}]^{s(d)}$  the column vector

$$\overrightarrow{[\underline{X}]}_d := [1 \quad X_1 \quad X_2 \quad \dots \quad X_n \quad X_1^2 \quad X_1 X_2 \quad \dots \quad \dots \quad X_n^d]^*$$

consisting of these monomials ordered first with respect to the degree and then lexicographic.

The following proposition shows how to find elements of degree at most  $d + e$  (represented by a matrix  $W$ ) in the real radical  $\mathcal{I} := \sqrt[q]{\mathbb{R}}$  of the ideal generated by a polynomial  $q \in \mathbb{R}[\underline{X}]_{2d}$  (represented by a symmetric matrix  $U$ , i.e.,  $q = \overrightarrow{[\underline{X}]}_d^* U \overrightarrow{[\underline{X}]}_d$ ). We will later use it with  $d = 1$  and  $e = 2$  since  $q$  will be quadratic and we will need cubic elements in  $\mathcal{I}$ . Note that

$$U \succeq W^* W \iff \begin{bmatrix} I & W \\ W^* & U \end{bmatrix} \succeq 0$$

by the method of Schur complements.

**Proposition 3.4.1.** *Let  $d, e \in \mathbb{N}_0$ ,  $\mathcal{I}$  a real radical ideal of  $\mathbb{R}[\underline{X}]$  and  $U \in S\mathbb{R}^{s(d) \times s(d)}$  such that  $\overrightarrow{[\underline{X}]}_d^* U \overrightarrow{[\underline{X}]}_d \in \mathcal{I}$ . Suppose  $W \in \mathbb{R}^{s(e) \times s(d)}$  with  $U \succeq W^* W$ . Then  $\overrightarrow{[\underline{X}]}_e^* W \overrightarrow{[\underline{X}]}_d \in \mathcal{I}$ .*

*Proof.* Since  $U - W^* W$  is positive semidefinite, we find  $B \in \mathbb{R}^{s(d) \times s(d)}$  with  $U - W^* W = B^* B$ . Now let  $p_i \in \mathbb{R}[\underline{X}]$  denote the  $i$ -th entry of  $W \overrightarrow{[\underline{X}]}_d$  and  $q_j$  the  $j$ -th entry of  $B \overrightarrow{[\underline{X}]}_d$ . From

$$\begin{aligned} p_1^2 + \dots + p_{s(e)}^2 + q_1^2 + \dots + q_{s(d)}^2 &= (W \overrightarrow{[\underline{X}]}_d)^* W \overrightarrow{[\underline{X}]}_d + (B \overrightarrow{[\underline{X}]}_d)^* B \overrightarrow{[\underline{X}]}_d \\ &= \overrightarrow{[\underline{X}]}_d^* (W^* W + B^* B) \overrightarrow{[\underline{X}]}_d = \overrightarrow{[\underline{X}]}_d^* U \overrightarrow{[\underline{X}]}_d \in \mathcal{I} \end{aligned}$$

it follows that  $p_1, \dots, p_{s(e)} \in \mathcal{I}$  since  $\mathcal{I}$  is real radical. Now

$$\overrightarrow{[\underline{X}]}_e^* W \overrightarrow{[\underline{X}]}_d = \overrightarrow{[\underline{X}]}_e^* [p_1 \dots p_{s(e)}]^* = [p_1 \dots p_{s(e)}] \overrightarrow{[\underline{X}]}_e \in \mathcal{I}$$

since  $\mathcal{I}$  is an ideal. ■

The following lemma is a weak converse to Proposition 3.4.1. Its proof relies heavily on the fact that only linear and quadratic polynomials are involved.

**Lemma 3.4.2.** *Suppose  $\ell_1, \dots, \ell_t \in \mathbb{R}[\underline{X}]_1$  and  $q_1, \dots, q_t \in \mathbb{R}[\underline{X}]_2$ . Let  $U \in S\mathbb{R}^{s(1) \times s(1)}$  be such that*

$$\overrightarrow{[\underline{X}]}_1^* U \overrightarrow{[\underline{X}]}_1 = \ell_1^2 + \dots + \ell_t^2.$$

*Then there exists  $\lambda > 0$  and  $W \in \mathbb{R}^{s(2) \times s(1)}$  such that  $\lambda U \succeq W^* W$  and*

$$\overrightarrow{[\underline{X}]}_2^* W \overrightarrow{[\underline{X}]}_1 = \ell_1 q_1 + \dots + \ell_t q_t.$$

*Proof.* Suppose that at least one  $q_i \neq 0$  (otherwise take  $W = 0$ ). Choose column vectors  $c_i \in \mathbb{R}^{s(2)}$  such that  $c_i^* \overrightarrow{[\underline{X}]}_2 = q_i$ . Now let  $W \in \mathbb{R}^{s(2) \times s(1)}$  be the matrix defined by  $W \overrightarrow{[\underline{X}]}_1 = \sum_{i=1}^t \ell_i c_i$  so that

$$\overrightarrow{[\underline{X}]}_2^* W \overrightarrow{[\underline{X}]}_1 = \sum_{i=1}^t \ell_i \overrightarrow{[\underline{X}]}_2^* c_i = \sum_{i=1}^t \ell_i c_i^* \overrightarrow{[\underline{X}]}_2 = \sum_{i=1}^t \ell_i q_i.$$

Moreover, we get

$$\overrightarrow{[X]}_1^* W^* W \overrightarrow{[X]}_1 = (W \overrightarrow{[X]}_1)^* W \overrightarrow{[X]}_1 = \sum_{i,j=1}^t (\ell_i c_i)^* (\ell_j c_j)$$

and therefore for all  $x \in \mathbb{R}^n$

$$\begin{aligned} \begin{bmatrix} 1 & x_1 & \dots & x_n \end{bmatrix} W^* W \begin{bmatrix} 1 \\ x_1 \\ \vdots \\ x_n \end{bmatrix} &= \sum_{i,j=1}^t (\ell_i(x) c_i)^* (\ell_j(x) c_j) \\ &\leq \frac{1}{2} \sum_{i,j=1}^t ((\ell_i(x) c_i)^* (\ell_i(x) c_i) + (\ell_j(x) c_j)^* (\ell_j(x) c_j)) \\ &\leq t \sum_{i=1}^t (\ell_i(x) c_i)^* (\ell_i(x) c_i) \leq \lambda \sum_{i=1}^t \ell_i(x)^2, \end{aligned}$$

where we set  $\lambda := t \sum_{i=1}^t c_i^* c_i > 0$ . Therefore

$$\begin{bmatrix} 1 & x_1 & \dots & x_n \end{bmatrix} (\lambda U - W^* W) \begin{bmatrix} 1 & x_1 & \dots & x_n \end{bmatrix}^* \geq 0$$

for all  $x \in \mathbb{R}^n$ . By homogeneity and continuity this implies  $y^* (\lambda U - W^* W) y \geq 0$  for all  $y \in \mathbb{R}^{s(1)}$ , i.e.,  $\lambda U \succeq W^* W$ .  $\blacksquare$

**3.5. The sums of squares dual of an SDP.** Given an SDP of the form (P) described in Subsection 3.1, the following is what we call its *sums of squares dual*:

$$\begin{aligned} &\text{maximize } a \\ &\text{subject to } S \in S\mathbb{R}^{m \times m}, S \succeq 0, a \in \mathbb{R} \\ &\quad S_1, \dots, S_n \in S\mathbb{R}[\overrightarrow{[X]}]^{m \times m} \text{ quadratic sos-matrices} \\ &\quad U_1, \dots, U_n \in S\mathbb{R}^{s(1) \times s(1)} \\ &\quad W_1, \dots, W_n \in \mathbb{R}^{s(2) \times s(1)} \\ &\quad \overrightarrow{[X]}_1^* U_i \overrightarrow{[X]}_1 + \overrightarrow{[X]}_2^* W_{i-1} \overrightarrow{[X]}_1 + \text{tr}(LS_i) = 0 \quad (i \in \{1, \dots, n\}) \\ &\quad U_i \succeq W_i^* W_i \quad (i \in \{1, \dots, n\}) \\ &\quad \ell - a + \overrightarrow{[X]}_2^* W_n \overrightarrow{[X]}_1 - \text{tr}(LS) = 0. \end{aligned}$$

Just like Ramana's extended Lagrange-Slater dual [Ra] it can be written down in polynomial time (and hence has polynomial size) in the bit size of the primal (assuming the latter has rational coefficients) and it guarantees that strong duality (i.e., weak duality, zero gap and dual attainment, see Subsection 3.1) always holds. Similarly, the facial reduction [BW, TW] gives rise to a good duality theory of SDP. We refer the reader to [Pat] for a unified treatment of these two constructions.

As mentioned at the beginning of Section 3, the quadratic sos-matrices can easily be modeled by SDP constraints using the Gram matrix method, and the polynomial identities can be written as linear equations by comparing coefficients.

The  $S_i$  serve to produce negated quadratic sos-polynomials vanishing on  $S_L$  (cf. Proposition 3.2.1) which are captured by the matrices  $U_i$ . From this, cubics vanishing on  $S_L$  are produced (cf. Subsection 3.4) and represented by the matrices  $W_i$ . These cubics serve to implement the congruence modulo the ideals from (13) and (14). The whole procedure is iterated  $n$  times.

Just as Proposition 3.1.1 corresponds to the standard SDP duality, Theorem 3.5.2 translates into the strong duality for the sums of squares dual. Before we come to it, we need a folk lemma which is well-known (e.g. from the theory of Gröbner bases) and which we prove for the convenience of the reader.

**Lemma 3.5.1.** *Suppose  $d \in \mathbb{N}$ ,  $f \in \mathbb{R}[\underline{X}]_d$  and  $\ell_1, \dots, \ell_t \in \mathbb{R}[\underline{X}]_1$  are non-constant linear polynomials such that  $f \in (\ell_1, \dots, \ell_t)$ . Then there exist  $p_1, \dots, p_t \in \mathbb{R}[\underline{X}]_{d-1}$  such that  $f = p_1\ell_1 + \dots + p_t\ell_t$ .*

*Proof.* We proceed by induction on  $t \in \mathbb{N}_0$ . For  $t = 0$ , there is nothing to show. Now let  $t \in \mathbb{N}$  and suppose the lemma is already proved with  $t$  replaced by  $t - 1$ . We may assume that  $\ell_1 = X_1 - \ell$  with  $\ell \in \mathbb{R}[X_2, \dots, X_n]$  (otherwise permute the variables appropriately and scale  $\ell_1$ ). Write  $f = \sum_{\substack{|\alpha| \leq d \\ 1 \leq \alpha_1}} a_\alpha X^\alpha$  with  $a_\alpha \in \mathbb{R}$ . Setting  $g := f(\ell, X_2, \dots, X_n)$ , we have

$$f - g = \sum_{\substack{|\alpha| \leq d \\ 1 \leq \alpha_1}} a_\alpha (X_1^{\alpha_1} - \ell^{\alpha_1}) X_2^{\alpha_2} \dots X_n^{\alpha_n} = p_1(X_1 - \ell) = p_1\ell_1$$

where  $p_1 := \sum_{\substack{|\alpha| \leq d \\ 1 \leq \alpha_1}} a_\alpha \left( \sum_{i=0}^{\alpha_1-1} X_1^i \ell^{\alpha_1-1-i} \right) X_2^{\alpha_2} \dots X_n^{\alpha_n} \in \mathbb{R}[\underline{X}]_{d-1}$ . Moreover,  $g \in (\ell_2, \dots, \ell_t)$  and therefore  $g = p_2\ell_2 + \dots + p_t\ell_t$  for some  $p_2, \dots, p_t \in \mathbb{R}[\underline{X}]_{d-1}$  by induction hypothesis. Now

$$f = (f - g) + g = p_1\ell_1 + \dots + p_t\ell_t. \quad \blacksquare$$

**Theorem 3.5.2** (Sums of squares SDP duality). *Let  $L \in S\mathbb{R}[\underline{X}]^{m \times m}$  be a linear pencil and  $f \in \mathbb{R}[\underline{X}]_1$ . Then*

$$f \geq 0 \text{ on } S_L$$

*if and only if there exist quadratic sos-matrices  $S_1, \dots, S_n \in S\mathbb{R}[\underline{X}]^{m \times m}$ , matrices  $U_1, \dots, U_n \in S\mathbb{R}^{s(1) \times s(1)}$ ,  $W_1, \dots, W_n \in \mathbb{R}^{s(2) \times s(1)}$ ,  $S \in S\mathbb{R}_{\geq 0}^{m \times m}$  and  $c \in \mathbb{R}_{\geq 0}$  such that*

$$\overrightarrow{[\underline{X}]_1}^* U_i \overrightarrow{[\underline{X}]_1} + \overrightarrow{[\underline{X}]_2}^* W_{i-1} \overrightarrow{[\underline{X}]_1} + \text{tr}(LS_i) = 0 \quad (i \in \{1, \dots, n\}), \quad (18)$$

$$U_i \succeq W_i^* W_i \quad (i \in \{1, \dots, n\}), \quad (19)$$

$$f - c + \overrightarrow{[\underline{X}]_2}^* W_n \overrightarrow{[\underline{X}]_1} - \text{tr}(LS) = 0, \quad (20)$$

where  $W_0 := 0$ .

*Proof.* We first prove that existence of the above data implies  $f \geq 0$  on  $S_L$ . All we will use about the traces appearing in (18) and (20) is that they are polynomials nonnegative on  $S_L$ . Let  $\mathcal{I}$  denote the real radical ideal of all polynomials vanishing on  $S_L$ . It is clear that (20) gives  $f \geq 0$  on  $S_L$  if we show that  $\overrightarrow{[\underline{X}]_2} W_n \overrightarrow{[\underline{X}]_1} \in \mathcal{I}$ . In fact, we prove by induction that  $\overrightarrow{[\underline{X}]_2}^* W_i \overrightarrow{[\underline{X}]_1} \in \mathcal{I}$  for all  $i \in \{0, \dots, n\}$ .

The case  $i = 0$  is trivial since  $W_0 = 0$  by definition. Let  $i \in \{1, \dots, n\}$  be given and suppose that  $\overrightarrow{[\underline{X}]_2}^* W_{i-1} \overrightarrow{[\underline{X}]_1} \in \mathcal{I}$ . Then (18) shows  $\overrightarrow{[\underline{X}]_1}^* U_i \overrightarrow{[\underline{X}]_1} \leq 0$  on  $S_L$ . On the other hand (19) implies in particular  $U_i \succeq 0$  and therefore  $\overrightarrow{[\underline{X}]_1}^* U_i \overrightarrow{[\underline{X}]_1} \geq 0$  on  $S_L$ . Combining both,  $\overrightarrow{[\underline{X}]_1}^* U_i \overrightarrow{[\underline{X}]_1} \in \mathcal{I}$ . Now Proposition 3.4.1 implies  $\overrightarrow{[\underline{X}]_2}^* W_i \overrightarrow{[\underline{X}]_1} \in \mathcal{I}$  by (19). This ends the induction and shows  $f \geq 0$  on  $S_L$  as claimed.

Conversely, suppose now that  $L$  is infeasible. By Theorem 3.3.1 and Lemma 3.5.1, we can choose  $\ell_1, \dots, \ell_n \in \mathbb{R}[\underline{X}]_1$ , quadratic sos-matrices  $S'_1, \dots, S'_n \in S\mathbb{R}[\underline{X}]^{m \times m}$ ,  $S \in S\mathbb{R}_{\geq 0}^{m \times m}$  and  $q_{ij} \in \mathbb{R}[\underline{X}]_2$  ( $1 \leq j \leq i \leq n$ ) such that

$$\ell_1^2 + \dots + \ell_i^2 + \text{tr}(LS'_i) = \sum_{j=1}^{i-1} q_{(i-1)j} \ell_j \quad (i \in \{1, \dots, n\}) \quad \text{and} \quad (21)$$

$$f - c - \text{tr}(LS) = \sum_{j=1}^n q_{nj} \ell_j. \quad (22)$$

There are two little arguments involved in this: First, (13) can trivially be rewritten as  $\ell_1^2 + \dots + \ell_i^2 + \text{tr}(LS'_i) \in (\ell_1, \dots, \ell_{i-1})$  for  $i \in \{1, \dots, n\}$ . Second, in Lemma 3.5.1 the  $\ell_i$  are assumed to be non-constant but can be allowed to equal zero. But if

some  $\ell_i \neq 0$  is constant, then we may set  $\ell_{i+1} = \dots = \ell_n = 0$  and  $S'_{i+1} = \dots = S'_n = S = 0$ .

Now define  $U'_i \in S\mathbb{R}^{s(1) \times s(1)}$  by  $\overline{[X]}_1^* U'_i \overline{[X]}_1 = \ell_1^2 + \dots + \ell_i^2$  for  $i \in \{1, \dots, n\}$ . Using Lemma 3.4.2, we can then choose  $\lambda > 0$  and  $W'_1, \dots, W'_n \in \mathbb{R}^{s(2) \times s(1)}$  such that

$$\lambda U'_i \succeq W_i'^* W'_i \quad (23)$$

and  $\overline{[X]}_2^* W'_i \overline{[X]}_1 = -\sum_{j=1}^i q_{ij} \ell_j$  for  $i \in \{1, \dots, n\}$ . Setting  $W_n := W'_n$ , equation (22) becomes (20). Moreover, (21) can be rewritten as

$$\overline{[X]}_1^* U'_i \overline{[X]}_1 + \overline{[X]}_2^* W'_{i-1} \overline{[X]}_1 + \text{tr}(L S'_i) = 0 \quad (i \in \{1, \dots, n\}) \quad (24)$$

To cope with the problem that  $\lambda$  might be larger than 1 in (23), we look for  $\lambda_1, \dots, \lambda_n \in \mathbb{R}_{>0}$  such that  $U_i \succeq W_i^* W_i$  for all  $i \in \{1, \dots, n\}$  if we define  $U_i := \lambda_i U'_i$  and  $W_{i-1} := \lambda_i W'_{i-1}$  for all  $i \in \{1, \dots, n\}$  (in particular  $W_0 = W'_0 = 0$ ). With this choice, the desired linear matrix inequality (19) is now equivalent to  $\lambda_i U'_i \succeq \lambda_{i+1}^2 W_i'^* W'_i$  for  $i \in \{1, \dots, n-1\}$  and  $\lambda_n U'_n \succeq W_n'^2$ . Looking at (23), we therefore see that any choice of the  $\lambda_i$  satisfying  $\lambda_i \geq \lambda \lambda_{i+1}^2$  for  $i \in \{1, \dots, n-1\}$  and  $\lambda_n \geq \lambda$  ensures (19). Such a choice is clearly possible. Finally, equation (24) multiplied by  $\lambda_i$  yields (18) by setting  $S_i := \lambda_i S'_i$  for  $i \in \{1, \dots, n\}$ . ■

**3.6. The real radical and the quadratic module.** Let  $L$  be a linear pencil. In Definition 1.2.1, we have introduced the convex cone  $C_L \subseteq \mathbb{R}[\underline{X}]_1$  and the quadratic module  $M_L \subseteq \mathbb{R}[\underline{X}]$  associated to  $L$  consisting of polynomials which are obviously nonnegative of the spectrahedron  $S_L$ . In Theorem 2.2.5, we have shown that the quadratic module  $M_L$  can be used to certify infeasibility of  $L$ , in the sense that  $S_L = \emptyset$  implies  $-1 \in M_L$ . On the contrary, we have seen in Subsection 2.4, that the convex cone  $C_L$  is in general too small to detect infeasibility of  $L$  in this way.

In this subsection, we turn over to the more general question of certifying non-negativity of *arbitrary* linear polynomials on  $S_L$  (as opposed to just the constant polynomial  $-1$ ). The following example shows that  $M_L$  (and henceforth its subset  $C_L$ ) does not, in general, contain all linear polynomials nonnegative on  $S_L$ .

**Example 3.6.1.** Consider

$$L = \begin{bmatrix} 1 & X \\ X & 0 \end{bmatrix}.$$

Then  $S_L = \{0\}$ . Hence obviously  $X \geq 0$  on  $S_L$ . But it is easy to see that  $X \notin M_L$  [Za, Example 2].

Despite this example, Theorem 3.3.1 and its SDP-implementable version Theorem 3.5.2 do on the other hand yield algebraic certificates of linear polynomials nonnegative on  $S_L$ . These two theorems have the advantage of being very well-behaved with respect to complexity issues but have the drawback of their statements being somewhat technical. Leaving complexity issues aside, one can however come back to a nice algebraic characterization of linear polynomials nonnegative on  $S_L$ .

Given  $f \in \mathbb{R}[\underline{X}]_1$  with  $f \geq 0$  on  $S_L$ , the certificates in Theorems 3.3.1 and 3.5.2 for being nonnegative on  $S_L$  can actually be interpreted as certificates of  $f$  lying in the convex cone  $C_L + \sqrt{\text{supp } M_L}$  by means of Prestel's theory of semiorderings. Note that each element of  $C_L + \sqrt{\text{supp } M_L}$  is of course nonnegative on  $S_L$  since the elements of  $\sqrt{\text{supp } M_L}$  vanish on  $S_L$ .

Finally, this will allow us to come back to the quadratic module  $M_L$ . We will show that it contains each linear polynomial nonnegative on  $S_L$  after adding an *arbitrarily small positive constant*, see Corollary 3.6.7.

In this subsection, basic familiarity with real algebraic geometry as presented e.g. in [BCR, Ma, PD] is needed. The following proposition follows easily from Prestel's theory of semiorderings on a commutative ring, see for example [Sc, 1.4.6.1].

**Proposition 3.6.2.** *Let  $M$  be a quadratic module in  $\mathbb{R}[\underline{X}]$ . Then*

$$\sqrt{\text{supp } M} = \sqrt[r]{\text{supp } M} = \bigcap \{\text{supp } S \mid S \text{ semiordering of } \mathbb{R}[\underline{X}], M \subseteq S\}.$$

We explicitly extract the following consequence since this is exactly what is needed in the sequel.

**Lemma 3.6.3.** *Let  $M$  be a quadratic module in  $\mathbb{R}[\underline{X}]$ . Then*

$$(\sqrt{\text{supp } M} - M) \cap M \subseteq \sqrt{\text{supp } M}. \quad (25)$$

*Proof.* To prove (25), suppose  $p \in M$  can be written  $p = g - q$  with  $g \in \sqrt{\text{supp } M}$  and  $q \in M$ . By Proposition 3.6.2, we have to show that  $p \in \text{supp } S$  for each semiordering  $S$  of  $\mathbb{R}[\underline{X}]$  with  $M \subseteq S$ . But if such  $S$  is given, then  $g \in \text{supp } S$  and therefore  $p = g - q \in -S$  as well as  $p \in M \subseteq S$ . Hence  $p \in \text{supp } S$ . ■

Having this lemma at hand, we can now give a conceptual interpretation of the certificates appearing in Theorem 3.3.1, disregarding the complexity of the certificate.

**Proposition 3.6.4.** *If  $L \in S\mathbb{R}[\underline{X}]^{m \times m}$  is a linear pencil,  $f, \ell_1, \dots, \ell_n \in \mathbb{R}[\underline{X}]_1$ ,  $S_1, \dots, S_n \in S\mathbb{R}[\underline{X}]^{m \times m}$  are quadratic sos-matrices,  $S \in S\mathbb{R}_{\geq 0}^{m \times m}$  and  $c \in \mathbb{R}_{\geq 0}$  such that (13) and (14) hold, then  $f \in C_L + \sqrt{\text{supp } M_L}$ .*

*Proof.* Set  $\mathcal{I} := \sqrt{\text{supp } M_L}$ . It is clear that (14) gives  $f \in C_L + \mathcal{I}$  if we prove that  $\ell_i \in \mathcal{I}$  for all  $i \in \{1, \dots, n\}$ . Fix  $i \in \{1, \dots, n\}$  and assume by induction that  $\ell_1, \dots, \ell_{i-1} \in \mathcal{I}$ . Then (13) implies  $\ell_i^2 + \text{tr}(LS_i) \in \mathcal{I}$  and therefore  $\ell_i^2 \in (\mathcal{I} - M_L) \cap \sum \mathbb{R}[\underline{X}]^2 \subseteq (\mathcal{I} - M_L) \cap M_L \subseteq \mathcal{I}$  by (25). ■

We get the same interpretation for the certificates from Theorem 3.5.2.

**Proposition 3.6.5.** *If  $L \in S\mathbb{R}[\underline{X}]^{m \times m}$  is a linear pencil,  $f \in \mathbb{R}[\underline{X}]_1$ ,  $S_1, \dots, S_n \in S\mathbb{R}[\underline{X}]^{m \times m}$  are quadratic sos-matrices,  $U_1, \dots, U_n \in S\mathbb{R}^{s(1) \times s(1)}$ ,  $W_1, \dots, W_n \in \mathbb{R}^{s(2) \times s(1)}$ ,  $S \in S\mathbb{R}_{\geq 0}^{m \times m}$  and  $c \in \mathbb{R}_{\geq 0}$  such that (18), (19) and (19) hold, then  $f \in C_L + \sqrt{\text{supp } M_L}$ .*

*Proof.* Set  $\mathcal{I} := \sqrt{\text{supp } M_L}$ . It is clear that constraint (20) gives  $f \in C_L + \mathcal{I}$  if we show that  $\overline{[\underline{X}]}_2 W_n \overline{[\underline{X}]}_1 \in \mathcal{I}$ . In fact, we show by induction that  $\overline{[\underline{X}]}_2 W_i \overline{[\underline{X}]}_1 \in \mathcal{I}$  for all  $i \in \{0, \dots, n\}$ .

The case  $i = 0$  is trivial since  $W_0 = 0$  by definition. Let  $i \in \{1, \dots, n\}$  be given and suppose that we know already  $\overline{[\underline{X}]}_2^* W_{i-1} \overline{[\underline{X}]}_1 \in \mathcal{I}$ . Then (18) shows  $\overline{[\underline{X}]}_1^* U_i \overline{[\underline{X}]}_1 \in \mathcal{I} - M_L$ . On the other hand (19) implies in particular  $U_i \succeq 0$  and therefore  $\overline{[\underline{X}]}_1^* U_i \overline{[\underline{X}]}_1 \in \sum \mathbb{R}[\underline{X}]^2 \subseteq M_L$ . But then  $\overline{[\underline{X}]}_1^* U_i \overline{[\underline{X}]}_1 \in (\mathcal{I} - M_L) \cap M_L \subseteq \mathcal{I}$  by (25). Now (19) yields  $\overline{[\underline{X}]}_2^* W_i \overline{[\underline{X}]}_1 \in \mathcal{I}$  by Proposition 3.4.1 since  $\mathcal{I}$  is real radical by Proposition 3.6.2. This ends the induction. ■

The following corollary is now a generalization of Proposition 3.1.1 working also for low-dimensional  $S_L$  (note that  $\text{supp } M_L = (0)$  if  $S_L$  has non-empty interior).

**Corollary 3.6.6.** *Let  $L \in S\mathbb{R}[\underline{X}]^{m \times m}$  be a linear pencil. Then*

$$f \geq 0 \text{ on } S_L \iff f \in C_L + \sqrt{\text{supp } M_L}$$

for all  $f \in \mathbb{R}[\underline{X}]_1$ .

*Proof.* Combine either Theorem 3.3.1 with Proposition 3.6.4, or Theorem 3.5.2 with Proposition 3.6.5. ■

Now we come back to the quadratic module where we cannot avoid to add  $\varepsilon > 0$  in order to get a certificate as was shown in Example 3.6.1.

**Corollary 3.6.7.** *Let  $L \in S\mathbb{R}[X]^{m \times m}$  be a linear pencil. Then*

$$f \geq 0 \text{ on } S_L \iff \forall \varepsilon > 0 : f + \varepsilon \in M_L$$

for all  $f \in \mathbb{R}[X]_1$ .

*Proof.* To prove the non-trivial implication, let  $f \in \mathbb{R}[X]_1$  with  $f \geq 0$  on  $S_L$  be given. It suffices to show  $f + \varepsilon \in M_L$  for the special case  $\varepsilon = 1$  (otherwise replace  $f$  by  $\varepsilon f$  and divide by  $\varepsilon$ ). By Corollary 3.6.6, there exists  $g \in C_L$ ,  $p \in \mathbb{R}[X]$  and  $k \in \mathbb{N}$  such that  $f = g + p$  and  $p^k \in \mathcal{I} := \text{supp } M_L$ . Now  $f + \varepsilon = f + 1 = g + (f - g) + 1 = g + (p + 1)$  and it is enough to show that  $p + 1 \in M_L$ . This will follow from the fact that the image of  $p + 1$  is a square in the quotient ring  $\mathbb{R}[X]/\mathcal{I}$ . Indeed since the image of  $p$  in  $\mathbb{R}[X]/\mathcal{I}$  is nilpotent (in fact the image of  $p^k$  is zero), we can simply write down a square root of this element using the *finite* Taylor expansion at 1 of the square root function in 1 given by the binomial series:

$$p + 1 \equiv \left( \sum_{i=0}^{k-1} \binom{\frac{1}{2}}{i} p^i \right)^2 \pmod{\mathcal{I}}. \quad \blacksquare$$

#### 4. MATRICIAL SPECTRAHEDRA AND COMPLETE POSITIVITY

In this section we revisit some of the main results from [HKM], where we considered *noncommutative (matricial) relaxations* of linear matrix inequalities under the assumption of strict feasibility. The purpose of this section is twofold. First, in Subsection 4.2 we explain which of the results from [HKM] generalize to *weakly feasible* linear matrix inequalities. Second, in Subsection 4.3 we explain how our results from Section 3 pertain to (completely) positive maps under the absence of positive definite elements.

**4.1. Matricial relaxations of LMIs.** Suppose

$$L = A_0 + \sum_{i=1}^n X_i A_i$$

is a linear pencil of size  $m$ . Given  $Z = \text{col}(Z_1, \dots, Z_n) := \begin{bmatrix} Z_1 \\ \vdots \\ Z_n \end{bmatrix} \in (S\mathbb{R}^{d \times d})^n$ , the

*evaluation*  $L(Z)$  is defined as

$$L(Z) = I_d \otimes A_0 + \sum_{i=1}^n Z_i \otimes A_i \in S\mathbb{R}^{dm \times dm}.$$

The *matricial spectrahedron* of a linear pencil  $L$  is

$$\mathfrak{S}_L := \bigcup_{d \in \mathbb{N}} \{Z \in (S\mathbb{R}^{d \times d})^n \mid L(Z) \succeq 0\}.$$

Let

$$\mathfrak{S}_L(d) = \{Z \in (S\mathbb{R}^{d \times d})^n \mid L(Z) \succeq 0\} = \mathfrak{S}_L \cap (S\mathbb{R}^{d \times d})^n.$$

for  $d \in \mathbb{N}$ . The set  $\mathfrak{S}_L(1) \subseteq \mathbb{R}^n$  is the feasibility set of the linear matrix inequality  $L(x) \succeq 0$  and coincides with  $S_L$  as introduced above. We call  $\mathfrak{S}_L$  *bounded* if there is an  $N \in \mathbb{N}$  with operator norm  $\|Z\| \leq N$  for all  $Z \in \mathfrak{S}_L$ .

**Lemma 4.1.1.** *For matrices  $Z_1, \dots, Z_n$  of the same size, we have*

$$\|\text{col}(Z_1, \dots, Z_n)\|^2 = \left\| \sum_{i=1}^n Z_i^* Z_i \right\|.$$

*Proof.* Suppose  $Z_j \in \mathbb{R}^{d \times d}$ . Expand  $\text{col}(Z_1, \dots, Z_n) \in \mathbb{R}^{dn \times d}$  to a  $dn \times dn$  matrix  $\hat{Z}$  by adding zero columns. Then

$$\begin{aligned} \|\text{col}(Z_1, \dots, Z_n)\|^2 &= \|\hat{Z}\|^2 = \|\hat{Z}^* \hat{Z}\| \\ &= \left\| \begin{bmatrix} \sum_i Z_i^* Z_i & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \end{bmatrix} \right\| = \left\| \sum_i Z_i^* Z_i \right\|. \quad \blacksquare \end{aligned}$$

Given linear pencils  $L_1$  and  $L_2$ ,

$$L_j = A_{j,0} + \sum_{i=1}^n X_i A_{j,i} \in S\mathbb{R}[\underline{X}]_1^{m_j \times m_j}, \quad j = 1, 2, \quad (26)$$

we shall be interested in the following inclusion for matricial spectrahedra:

$$\mathfrak{S}_{L_1} \subseteq \mathfrak{S}_{L_2}. \quad (27)$$

In this case we say that  $L_1$  *matricially dominates*  $L_2$ . If  $L_2 \in \mathbb{R}[\underline{X}]_1$  then (27) is equivalent to  $S_{L_1} \subseteq S_{L_2}$ :

**Proposition 4.1.2.** *Let  $L$  be a linear pencil of size  $m$  and let  $\ell \in \mathbb{R}[\underline{X}]_1$ . Then*

$$\ell|_{S_L} \geq 0 \quad \iff \quad \ell|_{\mathfrak{S}_L} \succeq 0.$$

*Proof.* The implication  $\Leftarrow$  is obvious as  $\mathfrak{S}_L \supseteq S_L$ . For the converse assume  $\ell|_{\mathfrak{S}_L} \not\succeq 0$  and choose  $Z = (Z_1, \dots, Z_n) \in \mathfrak{S}_L$  with  $\ell(Z) \not\succeq 0$ . Let  $v$  be an eigenvector of  $\ell(Z)$  with negative eigenvalue. For  $v^* Z v = (v^* Z_1 v, \dots, v^* Z_n v) \in S_L$  we have

$$L(v^* Z v) = (I_m \otimes v)^* L(Z) (I_m \otimes v) \succeq 0,$$

and

$$0 > v^* \ell(Z) v = \ell(v^* Z v),$$

whence  $\ell|_{S_L} \not\succeq 0$ . ■

**Corollary 4.1.3.** *Let  $L$  be a linear pencil. Then*

$$\mathfrak{S}_L \text{ is bounded} \quad \iff \quad S_L = \mathfrak{S}_L(1) \text{ is bounded.}$$

**Corollary 4.1.4.** *For a linear pencil  $L$ ,  $\mathfrak{S}_L = \emptyset$  if and only if  $S_L = \emptyset$ .*

Since empty spectrahedra were thoroughly analyzed in previous sections, in the sequel we shall always *assume*  $S_L \neq \emptyset$ . Moreover, we assume that there is  $Z \in \mathfrak{S}_L$  with  $L(Z) \neq 0$ . Then, by compressing and translating, we ensure

**(Asmp)**  $0 \neq A_0 \succeq 0$ .

If the interior of  $S_L$  is non-empty, we could further reduce to strictly feasible linear pencils (cf. [HKM, Proposition 2.1]), but this is not the case we are interested in here.

**Lemma 4.1.5.** *If  $S_L$  is bounded, then  $A_0, \dots, A_n$  are linearly independent.*

*Proof.* This is easy; or see [HKM, Proposition 2.6(2)]. ■

We now introduce subspaces to be used in our considerations:

$$\mathcal{S}_j = \text{span}\{A_{j,i} \mid i = 0, \dots, n\} = \text{span}\{L_j(z) \mid z \in \mathbb{R}^n\} \subseteq S\mathbb{R}^{m_j \times m_j}.$$

We call  $\mathcal{S}_j = \mathcal{S}(L_j)$  the (nonunital) *operator system* associated to the linear pencil  $L_j$ . Conversely, to each linear subspace  $\mathcal{S} \subseteq S\mathbb{R}^{m \times m}$  we can associate a linear pencil  $L$  with  $\mathcal{S} = \mathcal{S}(L)$  by fixing a basis for  $\mathcal{S}$ . We can even enforce additional properties on  $L$ , see Proposition 4.3.2 below.

The key tool in studying inclusions of matricial spectrahedra as in (27) is the linear map  $\tau : \mathcal{S}(L_1) \rightarrow \mathcal{S}(L_2)$  we now define.



**Definition 4.1.6.** Let  $L_1, L_2$  be linear pencils as in (26). If  $A_{1,0}, \dots, A_{1,n}$  are linearly independent (e.g.  $S_{L_1}$  is bounded), we define the linear map

$$\tau : \mathcal{S}_1 \rightarrow \mathcal{S}_2, \quad A_{1,i} \mapsto A_{2,i} \quad (i = 0, \dots, n).$$

We shall soon see that, assuming (27),  $\tau$  has a property called complete positivity, which we now introduce. Let  $\mathcal{S}_j \subseteq S\mathbb{R}^{m_j \times m_j}$  be linear subspaces and  $\varphi : \mathcal{S}_1 \rightarrow \mathcal{S}_2$  a linear map. For  $d \in \mathbb{N}$ ,  $\varphi$  induces the map

$$\varphi_d = I_d \otimes \varphi : \mathbb{R}^{d \times d} \otimes \mathcal{S}_1 = \mathcal{S}_1^{d \times d} \rightarrow \mathcal{S}_2^{d \times d}, \quad M \otimes A \mapsto M \otimes \varphi(A),$$

called an *ampliation* of  $\varphi$ . Equivalently,

$$\varphi_d \left( \begin{bmatrix} T_{11} & \cdots & T_{1d} \\ \vdots & \ddots & \vdots \\ T_{d1} & \cdots & T_{dd} \end{bmatrix} \right) = \begin{bmatrix} \varphi(T_{11}) & \cdots & \varphi(T_{1d}) \\ \vdots & \ddots & \vdots \\ \varphi(T_{d1}) & \cdots & \varphi(T_{dd}) \end{bmatrix}$$

for  $T_{ij} \in \mathcal{S}_1$ . We say that  $\varphi$  is *d-positive* if  $\varphi_d$  is a positive map, i.e., maps positive semidefinite matrices into positive semidefinite matrices. If  $\varphi$  is *d-positive* for every  $d \in \mathbb{N}$ , then  $\varphi$  is *completely positive*.

**Remark 4.1.7.** If  $\mathcal{S}_1$  does not contain positive semidefinite matrices, then every linear map out of  $\mathcal{S}_1$  is completely positive. On the other hand, under the presence of positive definite elements there is a well established theory of completely positive maps, cf. [Pau]. In the absence of positive definite elements in  $\mathcal{S}_1$  the structure of completely positive maps out of  $\mathcal{S}_1$  seems to be more complicated, and our results from Section 3 describe (completely) positive maps  $\mathcal{S}_1 \rightarrow \mathbb{R}$  in this case; see Subsection 4.3 below.

**4.2. The map  $\tau$  is completely positive: LMI matricial domination.** The main result of this section is the equivalence of *d-positivity* of the map  $\tau$  from Definition 4.1.6 with the inclusion  $\mathfrak{S}_{L_1}(d) \subseteq \mathfrak{S}_{L_2}(d)$ . As a corollary,  $L_1$  matricially dominates  $L_2$  if and only if  $\tau$  is completely positive.

**Theorem 4.2.1** (cf. [HKM, Theorem 3.5]). *Let*

$$L_j = A_{j,0} + \sum_{i=1}^n X_i A_{j,i} \in S\mathbb{R}[\underline{X}]_1^{d_j \times d_j}, \quad j = 1, 2$$

*be linear pencils (with  $0 \neq A_{1,0} \succeq 0$ ), and assume the spectrahedron  $S_{L_1}$  is bounded. Let  $\tau$  be the linear map given in Definition 4.1.6.*

- (1)  $\tau$  is *d-positive* if and only if  $\mathfrak{S}_{L_1}(d) \subseteq \mathfrak{S}_{L_2}(d)$ ;
- (2)  $\tau$  is *completely positive* if and only if  $\mathfrak{S}_{L_1} \subseteq \mathfrak{S}_{L_2}$ .

We point out that in [HKM] this theorem has been given under an additional assumption: the linear pencil  $L_1$  was assumed to be strictly feasible.

To prove the theorem we start with a lemma.

**Lemma 4.2.2.** *Let  $L = A_0 + \sum_{i=1}^n X_i A_i$  be a linear pencil of size  $m$  with  $0 \neq A_0 \succeq 0$  defining a bounded spectrahedron  $S_L$ . Then:*

- (1) if  $\Lambda \in \mathbb{R}^{d \times d}$  and  $Z \in (S\mathbb{R}^{d \times d})^n$ , and if

$$S := \Lambda \otimes A_0 + \sum_{i=1}^n Z_i \otimes A_i \tag{28}$$

*is symmetric, then  $\Lambda = \Lambda^*$ ;*

- (2) if  $S \succeq 0$ , then  $\Lambda \succeq 0$ .

*Proof.* To prove item (1), suppose  $S$  is symmetric. Then  $0 = S - S^* = (\Lambda - \Lambda^*) \otimes A_0$ . Since  $A_0 \neq 0$ ,  $\Lambda = \Lambda^*$ .

For (2), if  $\Lambda \not\geq 0$ , then there is a unit vector  $v$  such that  $v^* \Lambda v < 0$ . Consider the orthogonal projection  $P \in S\mathbb{R}^{dm \times dm}$  from  $\mathbb{R}^d \otimes \mathbb{R}^m$  onto  $\mathbb{R}v \otimes \mathbb{R}^m$ , and let  $Y = ((v^* Z_i v) P_v)_{i=1}^n \in (S\mathbb{R}^{d \times d})^n$ . Here  $P_v \in S\mathbb{R}^{d \times d}$  is the orthogonal projection from  $\mathbb{R}^d$  onto  $\mathbb{R}v$ . Note that  $P = P_v \otimes I_m$ . Then the compression

$$PSP = P(\Lambda \otimes A_0 + \sum_{i=1}^n Z_i \otimes A_i)P = (v^* \Lambda v) P_v \otimes A_0 + \sum_{i=1}^n Y_i \otimes A_i \succeq 0,$$

which says that  $0 \neq \sum_{i=1}^n Y_i \otimes A_i \succeq 0$  since  $0 \neq A_0 \succeq 0$  and  $v^* \Lambda v < 0$ . This implies  $0 \neq tY \in \mathfrak{S}_L$  for all  $t > 0$ ; contrary to  $\mathfrak{S}_L$  being bounded.  $\blacksquare$

*Proof of Theorem 4.2.1.* In both of the statements, the direction ( $\Rightarrow$ ) is obvious. We focus on the converses.

Fix  $d \in \mathbb{N}$ . Suppose  $S \in \mathcal{S}_1^{d \times d}$  is positive semidefinite. Then it is of the form (28) for some  $\Lambda \in \mathbb{R}^{d \times d}$  and  $Z \in (S\mathbb{R}^{d \times d})^d$ :

$$S = \Lambda \otimes A_{1,0} + \sum_{i=1}^n Z_i \otimes A_{1,i} \succeq 0.$$

By Lemma 4.2.2,  $\Gamma \succeq 0$ . If we replace  $\Lambda$  by  $\Lambda + \varepsilon I$  for some  $\varepsilon > 0$ , the resulting  $S = S_\varepsilon$  is still in  $\mathcal{S}_1^{d \times d}$ , so without loss of generality we may assume  $\Lambda \succ 0$ . Hence,

$$(\Lambda^{-\frac{1}{2}} \otimes I)S(\Lambda^{-\frac{1}{2}} \otimes I) = I \otimes A_{1,0} + \sum_{i=1}^n (\Lambda^{-\frac{1}{2}} Z_i \Lambda^{-\frac{1}{2}}) \otimes A_{1,i} \succeq 0.$$

Since  $\mathfrak{S}_1(d) \subseteq \mathfrak{S}_2(d)$ , this implies

$$I \otimes A_{2,0} + \sum_{i=1}^n (\Lambda^{-\frac{1}{2}} Z_i \Lambda^{-\frac{1}{2}}) \otimes A_{2,i} \succeq 0.$$

Multiplying on the left and right by  $\Lambda^{\frac{1}{2}} \otimes I$  shows

$$\tau(S_\varepsilon) = \Lambda \otimes A_{2,0} + \sum_{i=1}^n Z_i \otimes A_{2,i} \succeq 0.$$

An approximation argument now implies that if  $S \succeq 0$ , then  $\tau(S) \succeq 0$  and hence  $\tau$  is  $d$ -positive proving (1). Item (2) follows immediately.  $\blacksquare$

We next use the complete positivity of  $\tau$  under the assumption of LMI matrix domination to give an algebraic characterization of linear pencils  $L_1$  and  $L_2$  producing an inclusion  $\mathfrak{S}_{L_1} \subseteq \mathfrak{S}_{L_2}$ .

**Corollary 4.2.3** (Linear Positivstellensatz, cf. [HKM, Theorem 1.1]). *Let  $L_1$  and  $L_2$  be linear pencils of sizes  $m_1$  and  $m_2$ , respectively. Assume that  $S_{L_1}$  is bounded and  $\mathcal{S}_1$  contains a positive definite matrix (e.g.  $L_1$  is strictly feasible). Then the following are equivalent:*

- (i)  $\mathfrak{S}_{L_1} \subseteq \mathfrak{S}_{L_2}$ ;
- (ii) there exist  $\mu \in \mathbb{N}$  and  $V \in \mathbb{R}^{\mu m_1 \times m_2}$  such that

$$L_2 = V^*(I_\mu \otimes L_1)V. \quad (29)$$

Before turning to the proof of the corollary, we pause for a remark.

**Remark 4.2.4.** Equation (29) can be equivalently written as

$$L_2 = \sum_{j=1}^{\mu} V_j^* L_1 V_j, \quad (30)$$

where  $V_j \in \mathbb{R}^{m_1 \times m_2}$  and  $V = \text{col}(V_1, \dots, V_\mu)$ . Moreover,  $\mu$  can be uniformly bounded by Choi's characterization [Pau, Proposition 4.7] of completely positive maps between matrix algebras. In fact,  $\mu \leq m_1 m_2$ .

*Proof of Corollary 4.2.3.* The proof of the nontrivial implication (i)  $\Rightarrow$  (ii) proceeds as follows. First invoke Arveson's extension theorem [Pau, Theorem 7.5] to extend  $\tau$  from Definition 4.1.6 to a completely positive map  $\psi : \mathbb{R}^{m_1 \times m_1} \rightarrow \mathbb{R}^{m_2 \times m_2}$ . (*Caution:* this is where the existence of a positive definite matrix in  $\mathcal{S}_1$  is used.) Then apply the Stinespring representation theorem [Pau, Theorem 4.1] to obtain

$$\psi(a) = V^* \pi(a) V, \quad a \in \mathbb{R}^{m_1 \times m_1} \quad (31)$$

for some unital  $*$ -representation  $\pi : \mathbb{R}^{m_1 \times m_1} \rightarrow \mathbb{R}^{m_3 \times m_3}$  and  $V : \mathbb{R}^{m_1} \rightarrow \mathbb{R}^{m_3}$ . By dissecting the proof (or see [HKM, pp. 12-14]) we see that  $\pi$  is (unitarily equivalent to) a multiple of the identity representation, i.e.,  $\pi(a) = I_\mu \otimes a$  for some  $\mu \in \mathbb{N}$  and all  $a \in \mathbb{R}^{m_1 \times m_1}$ . Hence (31) implies (29).  $\blacksquare$

### 4.3. Examples and concluding remarks.

**Example 4.3.1.** Let us revisit Example 3.6.1 to show that the existence of a positive definite matrix in  $\mathcal{S}_1$  is needed for Corollary 4.2.3 to hold. Consider

$$L_1 = \begin{bmatrix} 1 & X \\ X & 0 \end{bmatrix}, \quad L_2 = X.$$

Then  $S_{L_1} = \{0\}$  and  $\mathfrak{S}_{L_1} = \{0\}$ . Hence obviously  $\mathfrak{S}_{L_1} \subseteq \mathfrak{S}_{L_2}$ . However,  $L_2$  does not admit a representation of the form (30). Indeed, such a certificate is equivalent to  $X \in C_{L_1}$ ; but it was already pointed out that  $X$  is not even a member of  $M_{L_1}$ .

We finish this paper by showing how our Theorem 3.3.1 (cf. Theorem 3.5.2) pertains to positive maps  $\mathcal{S} \rightarrow \mathbb{R}$  for a linear subspace  $\mathcal{S} \subseteq S\mathbb{R}^{m \times m}$ . For this we start by associating to  $\mathcal{S}$  a linear pencil  $L$  with  $\mathcal{S}(L) = \mathcal{S}$  and bounded  $S_L$ .

**Proposition 4.3.2.** *Let  $\mathcal{S} \subseteq S\mathbb{R}^{m \times m}$  be a linear subspace of dimension  $n+1 \geq 2$  containing a nonzero positive semidefinite matrix. Then there exist  $A_0, A_1, \dots, A_n \in \mathcal{S}$  such that*

- (1)  $A_0 \succeq 0$ ;
- (2)  $\mathcal{S} = \text{span}\{A_0, \dots, A_n\}$ ;
- (3) for the linear pencil  $L := A_0 + X_1 A_1 + \dots + X_n A_n$  the spectrahedron  $S_L$  is bounded.

*Proof.* We will use the trace inner product

$$(A, B) \mapsto \langle A, B \rangle := \text{tr}(AB)$$

on  $S\mathbb{R}^{m \times m}$ .

Let  $A_0$  be a maximum rank positive semidefinite matrix in  $\mathcal{S}$ . Without loss of generality we may assume

$$A_0 = \begin{bmatrix} I_s & 0 \\ 0 & 0_{m-s} \end{bmatrix}$$

for some  $1 \leq s \leq m$ .

**Claim.** If for some  $A_1 = \begin{bmatrix} A_{11} & A_{12} \\ A_{12}^* & A_{22} \end{bmatrix} \in \mathcal{S}$  with  $A_{11} \in S\mathbb{R}^{s \times s}$  we have  $\langle A_0, A_1 \rangle = 0$  and

$$A_0 + \lambda A_1 = \begin{bmatrix} I_s + \lambda A_{11} & \lambda A_{12} \\ \lambda A_{12}^* & \lambda A_{22} \end{bmatrix} \succeq 0 \quad \text{for all } \lambda \in \mathbb{R}_{\geq 0}, \quad (32)$$

then  $A_1 = 0$ .

*Explanation.* Since  $\langle A_0, A_1 \rangle = 0$ ,  $\text{tr}(A_{11}) = 0$ . This means that either  $A_{11} = 0$  or  $A_{11}$  has both positive and negative eigenvalues. In the latter case, fix an eigenvalue

$\mu < 0$  of  $A_{11}$ . Then for every  $\lambda \in \mathbb{R}$  with  $\lambda > -\mu^{-1} > 0$ , we have that  $I_s + \lambda A_{11} \not\geq 0$ , contradicting (32). So  $A_{11} = 0$ . If  $s = m$  we are done. Hence assume  $s < m$ .

Now

$$A_0 + \lambda A_1 = \begin{bmatrix} I_s & \lambda A_{12} \\ \lambda A_{12}^* & \lambda A_{22} \end{bmatrix} \succeq 0 \quad (33)$$

for all  $\lambda \in \mathbb{R}_{\geq 0}$ . Using Schur complements, (33) is equivalent to

$$\lambda A_{22} - \lambda^2 A_{12}^* A_{12} \succeq 0.$$

Hence  $A_{22} - \lambda A_{12}^* A_{12} \succeq 0$  for all  $\lambda \in \mathbb{R}_{\geq 0}$ . Equivalently,  $A_{12} = 0$  and  $A_{22} \succeq 0$ . If  $A_{22} \neq 0$ , then  $0 \preceq A_0 + A_1 \in \mathcal{S}$ , and

$$s = \text{rank}(A_0) < \text{rank}(A_0 + A_1),$$

contradicting the maximality of the rank of  $A_0$ .  $\square$

Take an arbitrary orthogonal basis  $A_0, A_1, \dots, A_n$  of  $\mathcal{S}$  containing  $A_0$ , and let  $L := A_0 + X_1 A_1 + \dots + X_n A_n$ . We claim that  $S_L$  is bounded.

Assume otherwise. Then there exists a sequence  $(x^{(k)})_k$  in  $\mathbb{R}^n$  such that  $\|x^{(k)}\| = 1$  for all  $k$ , and an increasing sequence  $t_k \in \mathbb{R}_{>0}$  tending to  $\infty$  such that  $L(t_k x^{(k)}) \succeq 0$ . By convexity this implies  $t_k x^{(r)} \in S_L$  for all  $r \geq k$ . Without loss of generality we assume the sequence  $(x^{(k)})_k$  converges to a vector  $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ . Clearly,  $\|x\| = 1$ . For any  $t \in \mathbb{R}_{\geq 0}$ ,  $t x^{(k)} \rightarrow t x$ , and for  $k$  big enough,  $t x^{(k)} \in S_L$  by convexity. So  $x$  satisfies  $L(t x) \succeq 0$  for all  $t \in \mathbb{R}_{\geq 0}$ . In other words,

$$A_0 + t(x_1 A_1 + \dots + x_n A_n) \geq 0$$

for all  $t \in \mathbb{R}_{\geq 0}$ . But now the claim implies  $x_1 A_1 + \dots + x_n A_n = 0$ , contradicting the linear independence of the  $A_j$ .  $\blacksquare$

**Lemma 4.3.3.** *Suppose  $\mathcal{S} \subseteq S\mathbb{R}^{m \times m}$  is a linear subspace and  $\varphi : \mathcal{S} \rightarrow \mathbb{R}$  is a positive map. Then  $\varphi$  is completely positive.*

*Proof.* This is well-known and easy, cf. [Pau, Proposition 3.8 or Theorem 3.9].  $\blacksquare$

As seen in Example 4.3.1 there does not exist a clean linear certificate for LMI matricial domination (or, equivalently, complete positivity). However, our Theorem 3.3.1 yields a nonlinear algebraic certificate for LMI (matricial) domination  $S_L \subseteq S_f$  in the case  $f$  is a size 1 linear pencil, i.e.,  $f \in \mathbb{R}[X]_1$ . Equivalently, Theorem 3.3.1 can be used to describe (completely) positive maps  $\tau : \mathcal{S} \rightarrow \mathbb{R}^{m \times m}$ , where  $\mathcal{S} \subseteq S\mathbb{R}^{m \times m}$  is a(ny) linear subspace.

Indeed, suppose  $\mathcal{S}$  contains a nonzero positive semidefinite matrix (otherwise  $\tau$  is automatically completely positive) and apply Proposition 4.3.2 to obtain a basis  $A_0, \dots, A_n$  of  $\mathcal{S}$ , and the linear pencil

$$L := A_0 + X_1 A_1 + \dots + X_n A_n$$

of size  $m$  with bounded  $S_L$ . Define

$$f := \tau(A_0) + X_1 \tau(A_1) + \dots + X_n \tau(A_n) \in \mathbb{R}[X]_1.$$

Then, by Theorem 4.2.1 (here is where the boundedness of  $S_L$  is used),  $\tau$  is completely positive if and only if  $\mathfrak{S}_L \subseteq \mathfrak{S}_f$ . By Proposition 4.1.2 the latter is equivalent to  $S_L \subseteq S_f$ , i.e.,  $f|_{S_L} \geq 0$ , and this is a situation completely characterized by Theorem 3.3.1.

## REFERENCES

- [Ba1] A. Barvinok: A remark on the rank of positive semidefinite matrices subject to affine constraints, *Discrete Comput. Geom.* **25** (2001), no. 1, 23–31 [6](#)
- [Ba2] A. Barvinok: *A course in convexity*, Graduate Studies in Mathematics **54**, Amer. Math. Soc. 2002 [5](#), [6](#)
- [BCR] J. Bochnak, M. Coste, M.F. Roy: *Real algebraic geometry*, Springer-Verlag 1998 [17](#), [21](#)
- [BEFB] S. Boyd, L. El Ghaoui, E. Feron, V. Balakrishnan: *Linear matrix inequalities in system and control theory*, SIAM Studies in Applied Mathematics **15**, SIAM 1994 [2](#)
- [Bo] F. Bohnenblust: Joint positiveness of matrices, Technical report, California Institute of Technology 1948 [6](#)
- [BV] S. Boyd, L. Vandenberghe: Semidefinite programming, *SIAM Rev.* **38** (1996), no. 1, 49–95 [1](#)
- [BW] J.M. Borwein, H. Wolkowicz: Facial reduction for a cone-convex programming problem, *J. Austral. Math. Soc. Ser. A* **30** (1980/81), no. 3, 369–380 [2](#), [19](#)
- [dK] E. de Klerk: Aspects of semidefinite programming, Interior point algorithms and selected applications. *Applied Optimization*, **65**, Kluwer Academic Publishers, Dordrecht, 2002 [2](#)
- [Fa] J. Farkas: Theorie der einfachen Ungleichungen, *J. reine angew. Math.* **124** (1902), 1–27 [5](#)
- [GLR] I. Gohberg, P. Lancaster, L. Rodman: *Matrix polynomials*, Computer Science and Applied Mathematics, Academic Press 1982 [3](#)
- [GN] J. Gouveia, T. Netzer: Positive polynomials and projections of spectrahedra, preprint <http://arxiv.org/abs/0911.2750> [5](#)
- [Go] M. Goemans: Semidefinite programming in combinatorial optimization, Lectures on mathematical programming, *Math. Program.* **79** (1997), no. 1-3, Ser. B, 143–161 [2](#)
- [HKM] J.W. Helton, I. Klep, S. McCullough: The matricial relaxation of a linear matrix inequality, preprint <http://arxiv.org/abs/1003.0908> [3](#), [5](#), [9](#), [23](#), [24](#), [25](#), [26](#), [27](#)
- [HS] C.W.J. Hol, C.W. Scherer: Matrix sum-of-squares relaxations for robust semi-definite programs, *Math. Program.* **107** (2006), no. 1-2, Ser. B, 189–211 [9](#), [13](#)
- [Ko] M. Kojima: Sums of squares relaxations of polynomial semidefinite programs, Technical report, Tokyo Institute of Technology, 2003 [13](#)
- [KS] I. Klep, M. Schweighofer: Pure states, positive matrix polynomials and sums of Hermitian squares, *Indiana Univ. Math. J.* **59** (2010), no. 3, 857–874 [9](#)
- [Las] J.B. Lasserre: *Moments, positive polynomials and their applications*, Imperial College Press Optimization Series **1**, Imperial College Press 2010 [2](#)
- [Lau] M. Laurent: Sums of squares, moment matrices and optimization over polynomials, Emerging applications of algebraic geometry, 157–270, IMA Vol. Math. Appl. **149**, Springer, New York, 2009; updated version available at <http://homepages.cwi.nl/~monique/files/moment-ima-update-new.pdf> [2](#), [13](#)
- [Ma] M. Marshall: *Positive polynomials and sums of squares*, Mathematical Surveys and Monographs **146**, Amer. Math. Soc. 2008 [2](#), [3](#), [4](#), [8](#), [9](#), [13](#), [17](#), [21](#)
- [Ne] A. Nemirovski: Advances in convex optimization: conic programming, *International Congress of Mathematicians*, Vol. **I**, 413–444, Eur. Math. Soc. 2007 [1](#)
- [NN] Y. Nesterov, A. Nemirovskii: *Interior-point polynomial algorithms in convex programming*, SIAM Studies in Applied Mathematics **13**, SIAM 1994 [1](#), [2](#)
- [Par] P.A. Parrilo: *Structured Semidefinite Programs and Semialgebraic Geometry Methods in Robustness and Optimization*, Ph.D. thesis, California Institute of Technology 2000 [2](#)
- [Pat] G. Pataki: A Simple Derivation of a Facial Reduction Algorithm, and Extended Dual Systems, preprint <http://www.unc.edu/~pataki/papers/fr.pdf> [19](#)
- [Pau] V. Paulsen: *Completely bounded maps and operator algebras*, Cambridge Studies in Advanced Mathematics **78**, Cambridge University Press 2002 [25](#), [27](#), [28](#)
- [PD] A. Prestel, C.N. Delzell: *Positive polynomials. From Hilbert's 17th problem to real algebra*, Springer-Verlag 2001 [2](#), [3](#), [7](#), [8](#), [17](#), [21](#)
- [PS] V. Powers, C. Scheiderer: The moment problem for non-compact semialgebraic sets, *Adv. Geom.* **1** (2001), no. 1, 71–88 [14](#)
- [Pu] M. Putinar: Positive polynomials on compact semi-algebraic sets, *Indiana Univ. Math. J.* **42** (1993), no. 3, 969–984 [9](#)
- [Ra] M. Ramana: An exact duality theory for semidefinite programming and its complexity implications, *Math. Program.* **77** (1997), no. 2, Ser. B, 129–162 [2](#), [19](#)
- [Sc] C. Scheiderer: Positivity and sums of squares: a guide to recent results, Emerging applications of algebraic geometry, 271–324, IMA Vol. Math. Appl. **149**, Springer-Verlag 2009 [2](#), [4](#), [21](#)

- [SIG] R.E. Skelton, T. Iwasaki, K.M. Grigoriadis: *A unified algebraic approach to linear control design*, Taylor & Francis 1998 [2](#)
- [Sm] K. Schmüdgen: The  $K$ -moment problem for compact semi-algebraic sets, *Math. Ann.* **289** (1991), no. 2, 203–206 [8](#)
- [Sr] A. Schrijver: *Theory of linear and integer programming*, Wiley-Interscience Series in Discrete Mathematics, John Wiley & Sons, Ltd., Chichester, 1986 [1](#), [2](#)
- [St] J. Sturm: Theory and algorithms of semidefinite programming, High performance optimization, 1–194, *Appl. Optim.* **33**, Kluwer Acad. Publ. 2000 [5](#), [12](#)
- [To] M.J. Todd: Semidefinite optimization, *Acta Numer.* **10** (2001), 515–560 [1](#)
- [TW] L. Tuncel, H. Wolkowicz, Strong Duality and Minimal Representations for Cone Optimization, preprint  
<http://orion.math.uwaterloo.ca/~hwolkowi/henry/reports/regclosegap.pdf> [19](#)
- [WSV] H. Wolkowicz, R. Saigal, L. Vandenberghe (editors): *Handbook of semidefinite programming. Theory, algorithms, and applications*, Kluwer Acad. Publ. 2000 [1](#), [12](#)
- [Za] A. Zalar: A note on a matrix version of the Farkas lemma, preprint  
<http://arxiv.org/abs/1012.5420> [21](#)

IGOR KLEP, UNIVERZA V MARIBORU, FAKULTETA ZA NARAVOSLOVJE IN MATEMATIKO, KOROŠKA 160, SI-2000 MARIBOR, SLOVENIA, AND UNIVERZA V LJUBLJANI, FAKULTETA ZA MATEMATIKO IN FIZIKO, JADRANSKA 21, SI-1111 LJUBLJANA, SLOVENIA

*E-mail address:* igor.klep@mf.uni-lj.si

MARKUS SCHWEIGHOFER, UNIVERSITÄT KONSTANZ, FACHBEREICH MATHEMATIK UND STATISTIK, 78457 KONSTANZ, ALLEMAGNE

*E-mail address:* markus.schweighofer@uni-konstanz.de